



Hardness and Approximation for the Geodetic Set Problem in Some Graph Classes

Dibyayan Chakraborty¹, Florent Foucaud², Harmender Gahlawat^{1(✉)},
Subir Kumar Ghosh³, and Bodhayan Roy⁴

¹ Indian Statistical Institute, Kolkata, India
harmendergahlawat@gmail.com

² Univ. Bordeaux, Bordeaux INP, CNRS, LaBRI, UMR5800, 33400 Talence, France

³ Ramakrishna Mission Vivekananda Educational and Research Institute,
Kolkata, India

⁴ Indian Institute of Technology, Kharagpur, Kharagpur, India

Abstract. In this paper, we study the computational complexity of finding the *geodetic number* of graphs. A set of vertices S of a graph G is a *geodetic set* if any vertex of G lies in some shortest path between some pair of vertices from S . The MINIMUM GEODETIC SET (MGS) problem is to find a geodetic set with minimum cardinality. In this paper, we prove that solving MGS is NP-hard on planar graphs with a maximum degree six and line graphs. We also show that unless $P = NP$, there is no polynomial time algorithm to solve MGS with sublogarithmic approximation factor (in terms of the number of vertices) even on graphs with diameter 2. On the positive side, we give an $O(\sqrt[3]{n} \log n)$ -approximation algorithm for MGS on general graphs of order n . We also give a 3-approximation algorithm for MGS on solid grid graphs which are planar.

1 Introduction and Results

Suppose there is a city-road network (i.e. a graph) and a bus company wants to open bus terminals in some of the cities. The buses will go from one bus terminal to another (i.e. from one city to another) following the shortest route in the network. Finding the minimum number of bus terminals required so that any city belongs to some shortest route between some pair of bus terminals is equivalent to finding the *geodetic number* of the corresponding graph. Formally, an undirected simple graph G has vertex set $V(G)$ and edge set $E(G)$. For two vertices $u, v \in V(G)$, let $I(u, v)$ denote the set of all vertices in G that lie in some shortest path between u and v . A set of vertices S is a *geodetic set* if $\cup_{u, v \in S} I(u, v) = V(G)$. The *geodetic number*, denoted as $g(G)$, is the minimum integer k such that G has a geodetic set of cardinality k . Given a graph G , the MINIMUM GEODETIC SET (MGS) problem is to compute a geodetic set of G with minimum cardinality. In this paper, we shall study the computational complexity of MGS in various graph classes.

The notion of geodetic sets and geodetic number was introduced by Harary et al. [18]. The notion of geodetic number is closely related to convexity and convex hulls in graphs, which have applications in game theory, facility location, information retrieval, distributed computing and communication networks [2, 10, 15, 19, 22]. In 2002, Atici [1] proved that finding the geodetic number of arbitrary graphs is NP-hard. Later, Dourado et al. [8, 9] strengthened the above result to *bipartite* graphs, *chordal* graphs and *chordal bipartite* graphs. Recently, Bueno et al. [3] proved that MGS remains NP-hard even for *subcubic* graphs. On the positive side, polynomial time algorithms to solve MGS are known for *cographs* [8], *split* graphs [8], *ptolemaic* graphs [12], *outer planar* graphs [21] and *proper interval* graphs [11]. In this paper, we prove the following theorem.

Theorem 1. *MGS is NP-hard for planar graphs of maximum degree 6.*

Then we focus on *line* graphs. Given a graph G , the *line* graph of G , denoted by $L(G)$, is a graph such that each vertex of $L(G)$ represents an edge of G and two vertices of $L(G)$ are adjacent if and only if their corresponding edges share a common endpoint in G . A graph H is a *line graph* if $H \cong L(G)$ for some G . Some optimisation problems which are difficult to solve in general graphs admit polynomial time algorithms when the input is a line graph [14, 17]. We prove the following theorem.

Theorem 2. *MGS is NP-hard for line graphs.*

From a result of Dourado et al. [8], it follows that solving MGS is NP-hard even for graphs with diameter at most 4. On the other hand, solving MGS on graphs with diameter 1 is trivial (since those are exactly complete graphs). In this paper, we prove that unless $P = NP$, there is no polynomial time algorithm with sublogarithmic approximation factor for MGS even on graphs with diameter at most 2. A *universal vertex* of a graph is adjacent to all other vertices of the graph. We shall prove the following stronger theorem.

Theorem 3. *Unless $P = NP$, there is no polynomial time $o(\log n)$ -approximation algorithm for MGS even on graphs that have a universal vertex, where n is the number of vertices in the input graph.*

On the positive side, we show that a reduction to the MINIMUM RAINBOW SUBGRAPH OF MULTIGRAPH problem (defined in Sect. 3.1) gives the first sub-linear approximation algorithm for MGS on general graphs.

Theorem 4. *Given a graph, there is a polynomial-time $O(\sqrt[3]{n} \log n)$ -approximation algorithm for MGS where n is the number of vertices.*

Then we focus on *solid grid* graphs, an interesting subclass of planar graphs. A *grid embedding* of a graph is a collection of points with integer coordinates such that each point in the collection represents a vertex of the graph and two points are at a distance one if and only if the vertices they represent are adjacent

in the graph. A graph is a *grid* graph if it has a grid embedding. A graph is a *solid grid* graph if it has a grid embedding such that all interior faces have unit area. Approximation algorithms for optimisation problems like LONGEST PATH, LONGEST CYCLE, NODE-DISJOINT PATH etc. on grid graphs and solid grid graphs have been studied [4, 6, 20, 23, 25, 27]. In this paper, we prove the following theorem.

Theorem 5. *Given a solid grid graph, there is an $O(n)$ time 3-approximation algorithm for MGS, even if the grid embedding is not given as part of the input. Here n is the number of vertices in the input graph.*

Note that recognising solid grid graphs is NP-complete [16].

Organisation of the Paper: In Sect. 2, we prove the hardness results for planar graphs, line graphs and graphs with diameter 2. In Sect. 3, we present our approximation algorithms. Finally we draw our conclusions in Sect. 4.

2 Hardness Results

In Sect. 2.1, we prove that MGS is NP-hard for planar graphs with maximum degree 6 (Theorem 1). Then in Sect. 2.2 we prove that MGS is NP-hard for line graphs (Theorem 2). In Sect. 2.3 we prove the inapproximability result (Theorem 3).

2.1 NP-hardness on Planar Graphs

Given a graph G , a subset $S \subseteq V(G)$ is a dominating set of G if any vertex in $V(G) \setminus S$ has a neighbour in S . The problem MINIMUM DOMINATING SET (MDS) consists in computing a dominating set of an input graph G with minimum cardinality. To prove Theorem 1, we reduce the NP-complete MDS on subcubic planar graphs [13] to MGS on planar graphs with maximum degree 6.

Let us describe the reduction. From a subcubic planar graph G with a given planar embedding, we construct a graph $f(G)$ as follows. Each vertex v of G will be replaced by a *vertex gadget* G_v . This vertex gadget has vertex set $\{c^v, t_0^v, t_1^v, t_2^v\} \cup \{x_{i,j}^v, y_{i,j}^v, z_{i,j}^v \mid 0 \leq i < j \leq 2\}$. For simplicity we will consider that $x_{i,j}^v$ and $x_{j,i}^v$ refers to the same vertex (the same holds for $y_{i,j}^v$ and $y_{j,i}^v$, and for $z_{i,j}^v$ and $z_{j,i}^v$). There are no other vertices in $f(G)$. For the edges within G_v , vertex t_i^v (for $0 \leq i \leq 2$) is adjacent to vertices c^v , $x_{i,i+1}^v$, $y_{i,i+1}^v$, $x_{i-1,i}^v$, $y_{i-1,i}^v$ (indices taken modulo 3). Moreover, for each pair i, j with $0 \leq i < j \leq 2$, $x_{i,j}^v$ is adjacent to c^v and $y_{i,j}^v$, and $y_{i,j}^v$ is adjacent to $z_{i,j}^v$. We now describe the edges outside of the vertex-gadgets. They will depend on the embedding of G . We assume that the edges incident with any vertex v are labeled e_i^v with $0 \leq i < \deg_G(v)$, in such a way that the numbering increases counterclockwise around v with respect to the embedding (thus the edge vw will have two labels: e_i^v and e_j^w). Consider two vertices v and w that are adjacent in G , and let e_i^v and e_j^w be the two labels of edge vw in G . Then, t_i^v is adjacent to t_j^w , $y_{i,i+1}^v$ is

adjacent to $y_{j-1,j}^w$ and $y_{i-1,i}^v$ is adjacent to $y_{j+1,j}^w$ (indices are taken modulo the degree of the original vertex of G). It is clear that a planar embedding of $f(G)$ can easily be obtained from the planar embedding of G . Thus $f(G)$ is planar and has maximum degree 6. The construction is depicted in Fig. 1, where v and w are adjacent in G and the edge vw is labeled e_0^v and e_0^w .

We will show that G has a dominating set of size k if and only if $f(G)$ has a geodetic set of size $3|V(G)| + k$.




Fig. 1. Illustration of the reduction used in the proof of Theorem 1. Here, two vertex gadgets G_v, G_w are depicted, with v and w adjacent in G . Dashed lines represent potential edges to other vertex-gadgets.

Assume first that G has a dominating set D of size k . We construct a geodetic set S of $f(G)$ of size $3|V(G)| + k$ as follows. For each vertex v in G , we add the three vertices $z_{i,j}^v$ ($0 \leq i < j \leq 2$) of G_v to S . If v is in D , we also add vertex c^v to S .

Let us show that S is indeed a geodetic set. First, we observe that, in any vertex gadget G_v that is part of $f(G)$, the unique shortest path between two distinct vertices $z_{i,j}^v, z_{i',j'}^v$ has length 4 and goes through vertices $y_{i,j}^v, t_0^v$ and $y_{i',j'}^v$ (where $\{k\} = \{i, j\} \cap \{i', j'\}$). Thus, it only remains to show that vertices c^v and $x_{i,j}^v$ ($0 \leq i < j \leq 2$) belong to some shortest path of vertices of S . Assume that v is a vertex of G in D . The shortest paths between c^v and $z_{i,j}^v$ have length 3 and one of them goes through vertex $x_{i,j}^v$. Thus, all vertices of G_v belong to some shortest path between vertices of S . Now, consider a vertex w of G adjacent to v and let $z_{i,j}^w$ be the vertex of G_w that is farthest from c^v . The shortest paths between c^v and $z_{i,j}^w$ have length 6; one of them goes through vertices c^w and $x_{i,j}^w$; two others go through the two other vertices $x_{i',j'}^w$ and $x_{i'',j''}^w$. Thus, S is a geodetic set.

For the converse, assume we have a geodetic set S' of $f(G)$ of size $3|V(G)| + k$. We will show that G has a dominating set of size k . First of all, observe that all the $3|V(G)|$ vertices of type $z_{i,j}^v$ are necessarily in S' , since they have degree 1. As observed earlier, the shortest paths between those vertices already go through all vertices of type t_i^v and $y_{i,j}^v$. However, no other vertex lies on a shortest path between two such vertices: these shortest paths always go through the boundary




Fig. 2. (a) A triangle-free graph G and (b) the graph H_G .

6-cycle of the vertex-gadgets. Let S'_0 be the set of the remaining k vertices of S' . These vertices are there to cover the vertices of type c^v and $x_{i,j}^v$. We construct a subset D' of $V(G)$ as follows: D' contains those vertices v of G whose vertex-gadget G_v contains a vertex of S'_0 . We claim that D' is a dominating set of G . Suppose by contradiction that there is a vertex v of G such that neither G_v nor any of G_w (with w adjacent to v in G) contains any vertex of S'_0 . Here also, the shortest paths between vertices of S always go through the boundary 6-cycle of G_v and thus, they never include vertex c_v , a contradiction. Thus, D' is a dominating set of size k , and we are done.

2.2 NP-hardness on Line Graphs

In this section, we prove that MGS remains NP-hard on line graphs. For a graph G and edges $e, e' \in E(G)$, define $d(e, e') = 1$ if e, e' shares a vertex, and $d(e, e') = i > 1$ if e, e' do not share a vertex and e' shares a vertex with an edge e'' with $d(e, e'') = i - 1$. A path between two edges e, e' is defined in the usual way.

Observation 1. *A path between two edges e, e' of a graph G corresponds to a path between the vertices e and e' in $L(G)$.*

Given a graph G , a set $S \subseteq E(G)$ is a *line geodetic set* of G if every edge $e \in E(G) \setminus S$ belongs to some shortest path between some pair of edges $\{e, e'\} \subseteq S$. Observation 1 implies the following.

Observation 2. *A graph G has a line geodetic set of cardinality k if and only if $L(G)$ has a geodetic set of size k .*

We shall show (in Lemma 8) that finding a line geodetic set of a graph with minimum cardinality is NP-hard. Then Observation 2 shall imply that solving MGS on line graphs is NP-hard. For the above purpose we need the following definition. Given a graph G , a set $S \subseteq E(G)$ is a *good edge set* if for any edge $e \in E(G) \setminus S$, there are two edges $e', e'' \in S$ such that (i) e lies in some shortest path between e' and e'' , and (ii) $d(e', e'')$ is 2 or 3.

Lemma 6. *Computing a good edge set of a triangle-free graph with minimum cardinality is NP-hard.*

Proof. We shall reduce the NP-complete EDGE DOMINATING SET problem on triangle-free graphs [26] to the problem of computing a good edge set of a graph with minimum cardinality on triangle-free graphs. Given a graph G , a set $S \subseteq E(G)$ is an *edge dominating set* of G if any edge $e \in E(G) \setminus S$ shares a vertex with some edge in S . The EDGE DOMINATING SET problem is to compute an edge dominating set of G with minimum cardinality.

Let G be a triangle-free graph. For each vertex $v \in V(G)$, take a new edge $x_v y_v$. Construct a graph G^* whose vertex set is the union of $V(G)$ and the set $\{x_v, y_v\}_{v \in V(G)}$ and $E(G^*) = E(G) \cup \{vx_v\}_{v \in V(G)} \cup \{x_v y_v\}_{v \in V(G)}$. Notice that G^* is a triangle-free graph and we shall show that G has an edge dominating set of cardinality k if and only if G^* has a good edge set of cardinality $k + n$ where $n = |V(G)|$.

Let S be an edge dominating set of G . For each $v \in G$, let H_v be written as x_v, y_v, z_v . Notice that the set $S \cup \{x_v y_v\}_{v \in V(G)}$ forms a good edge set of G^* and has cardinality $k + n$. Let S' be a good edge set of G^* of size at most $k + n$. Notice that for each $v \in V(G)$, S' must contain the edge $x_v y_v$. Hence, the cardinality of the set $S' \cap E(G)$ is at most k . Moreover, for each $e \in E(G^*) \cap E(G)$, there is an edge $e' \in S'$ which is at distance 2 from e . As S' is a good edge set of G^* , any edge in $E(G) \setminus S'$ shares a vertex with some edge of S' . Hence $S' \cap E(G)$ is an edge dominating set of G of cardinality at most k . \square

For a triangle-free graph G , let H_G be the graph with $V(H_G) = V(G) \cup \{a, b, c, d\}$ and $E(H_G) = E(G) \cup \{ab, cd\} \cup E'$ where $E' = \{bv\}_{v \in V(G)} \cup \{cv\}_{v \in V(G)}$. See Fig. 2(a) and (b) for an example. We prove the following proposition.

Lemma 7. *For a triangle-free graph G , there is a line geodetic set Q of H_G with minimum cardinality such that $Q \cap E' = \emptyset$.*

Proof. For a set $S \subseteq E(H_G)$, an edge $f \in S$ covers an edge $e \in E(H_G)$, if there is another edge $f' \in S$ such that e lies in the shortest path between f and f' . Notice that the edges $\{ab, cd\}$ lie in any line geodetic set of H_G and all edges in E' are covered by ab and cd . First we prove the following claims.

Claim 1. *Let Q be a line geodetic set of H_G and $e \in E' \cap Q$. If e does not cover any edge of $E(G)$, then $Q \setminus \{e\}$ is a line geodetic set of H_G .*

The proof of the above claim follows from the fact that all edges in $E' \cup \{ab, cd\}$ are covered by ab and cd .

Claim 2. *Let Q be a line geodetic set of H_G and $e \in E' \cap Q$. There is another edge $e' \in E(G) \setminus Q$ such that $(Q \cup \{e'\}) \setminus \{e\}$ is a line geodetic set of H_G .*

To prove the claim above, first we define the *eccentricity* of an edge $e \in E(H_G)$ to be the maximum shortest path distance between e and any other edge in $E(H_G)$. Notice that the eccentricity of any edge in E' is two and the eccentricity of any edge of $E(G)$ in H_G is at most three. Now remove all edges from $E' \cap Q$ which do not cover any edge of $E(G)$. By Claim 1, the resulting set, say Q' , is a line

geodetic set of H_G . Let e be an edge $Q' \cap E'$ and let $\{f_1, f_2, \dots, f_k\} \subseteq E(G) \setminus Q'$ be the set of edges covered by e . Since the eccentricity of e is two, there must exist e_1, e_2, \dots, e_k in Q' such that f_i has a common endpoint with both e and e_i for each $i \in \{1, 2, \dots, k\}$. Therefore the distance between e and e_i is two for each $i \in \{1, 2, \dots, k\}$. As G is triangle-free, $e_i \neq e_j$ for any $i, j \in \{1, 2, \dots, k\}$. Choose any edge $f_j \in \{f_1, f_2, \dots, f_k\}$. Observe that the distance between f_j and e_i is two when $i \neq j$. Therefore, for each $i \in \{1, 2, \dots, j-1, j+1, \dots, k\}$, the edge f_i lies in the shortest path between f_j and e_i . Therefore, $(Q' \cup \{f_j\}) \setminus \{e\}$ is a line geodetic set of H_G .

Given any line geodetic set P of H_G , we can use the arguments used in Claims 1 and 2 repeatedly on P to construct a line geodetic set Q of H_G such that $|Q| \leq |P|$ and $Q \cap E' = \emptyset$. Thus we have the proof. \square

Lemma 8. *Computing a line geodetic set of a graph with minimum cardinality is NP-hard.*

Proof. We shall reduce the NP-complete problem of computing a good edge set of a triangle-free graph with minimum cardinality (Lemma 6). Let G be a triangle-free graph. Construct the graph H_G as stated above (just before Lemma 7). The set E' is also defined as before. We shall show that a triangle-free graph G has a good edge set of cardinality k if and only if H_G has a line geodetic set of cardinality $k+2$.

Let P be a good edge set of G . Notice that, for each edge $e \in E(G)$, there are two edges $e', e'' \in P$ such that e belongs to a shortest path between e' and e'' in H_G . Also any edge of E' belongs to a shortest path between the edges ab and cd in H_G . Hence $P \cup \{ab, cd\}$ is a line geodetic set of H_G with cardinality $k+2$.

Let Q be a line geodetic set of H_G of size $k+2$. Notice that $\{ab, cd\} \subseteq Q$ and let $Q' = Q \setminus \{ab, cd\}$. Due to Lemma 7, we can assume that Q' does not contain any edge of E' . Let e be an edge in $E(G) \setminus Q'$ and let $e', e'' \in Q'$ such that e lies in some shortest path between e' and e'' in H_G . Since the distance between e' and e'' is at most three in H_G , it follows that Q' is a good edge set of G with cardinality k . \square

2.3 Inapproximability on Graphs with Diameter 2

Given a graph G , a set $S \subseteq V(G)$ is a *2-dominating set* of G if any vertex $w \in V(G) \setminus S$ has at least two neighbours in S . The 2-MDS problem is to compute a 2-dominating set of graphs with minimum cardinality. We shall use the following result.

Theorem 9 ([5, 7]). *Unless $P = NP$, there is no polynomial time $o(\log n)$ -approximation algorithm for the 2-MDS problem on triangle-free graphs.*

Lemma 10. *Let G be a triangle-free graph and G' be the graph obtained by adding an universal vertex v to G . A set S of vertices of G' is a geodetic set if and only if $S \setminus \{v\}$ is a 2-dominating set of G .*

Proof. Let S be a geodetic set of G' . Observe that for any vertex $u \in V(G) \setminus S$ there must exist vertices $u_1, u_2 \in S \setminus \{v\}$ such that $u \in I(u_1, u_2)$ and $u_1 u_2 \notin E(G)$. Hence, S is a 2-dominating set of G . Conversely, let S' be any 2-dominating set of G . For any vertex $u \in V(G) \setminus S'$ there exist $v, v' \in S'$ such that $uv, uv' \in E(G)$. Since G is triangle-free, v and v' are non-adjacent. Hence, $u \in I(v, v')$ and $S' \cup \{v\}$ is a geodetic set of G' . \square

The proof of Theorem 3 follows due to Lemma 10 and Theorem 9.

3 Approximation Algorithms

In Sects. 3.1 and 3.2 we present approximation algorithms for MGS on general graphs and solid grid graphs, respectively.

3.1 General Graphs

We will reduce the MINIMUM GEODETIC SET problem to the MINIMUM RAINBOW SUBGRAPH OF MULTIGRAPH (MRSM) problem. A subgraph H of an edge colored multigraph G is *colorful* if H contains at least one edge of each color. Given an edge colored multigraph G , the MRSM problem is to find a colorful subgraph of G of minimum cardinality. The following is a consequence of a result due to Tirodkar and Vishwanathan [24].

Theorem 11 ([24]). *Given an edge colored multigraph G , there is a polynomial time $O(\sqrt[3]{n} \log n)$ -approximation algorithm to solve the MRSM problem where $n = |V(G)|$.*

We note that Tirodkar and Vishwanathan [24] proved the above theorem for simple graphs only, but the proof works for multigraphs as well.

Given a graph G form an edge colored multigraph H_G as follows. The vertex set of H_G is the same as G . For each triplet of not necessarily distinct vertices (u, v, w) such that u lies in some shortest path between v and w , add an edge in H_G between v and w having the color u . Observe that, G has a geodetic set of cardinality k if and only if H_G has a colorful subgraph with k vertices. The proof of Theorem 4 follows from Theorem 11.

3.2 Solid Grid Graphs

In this section, we shall give a linear time 3-approximation algorithm for MGS on solid grid graphs. From now on G shall denote a solid grid graph and \mathcal{R} is a grid embedding of G where every interior face has unit area.

Let G be a solid grid graph. A path P of G is a *corner path* if (i) no vertex of P is a cut vertex, (ii) both end-vertices of P have degree 2, and (iii) all vertices except the end-vertices of P have degree 3. See Fig. 3(a) for an example. Observe that for a corner path P , either the x -coordinates of all vertices of P are the same or the y -coordinates of all vertices of P are the same. Moreover, all vertices of a corner path lie in the outer face of G . The next observation follows from the definition of corner path and the fact that G is a solid grid graph.

Observation 3. Let P be a corner path of G . Consider the set $Q = \{v \in V(G) : v \notin V(P), N(v) \cap P \neq \emptyset\}$. Then Q induces a path in G . Moreover, if the x -coordinates (resp. the y -coordinates) of all the vertices of P are the same, then the x -coordinates (resp. the y -coordinates) of all vertices in Q are the same.

We shall use Observation 3 to prove a lower bound on the geodetic number of G in terms of the number of corner paths of G .




Fig. 3. (a) The black and gray vertices are the vertices of the corner paths. The gray vertices indicate the corner vertices. (b) The gray vertices are vertices of the red path. Vertices in the shaded box form a rectangular block. (c) Example of a solid grid graph whose number of corner vertices is exactly three times the geodetic number.

Lemma 12. Any geodetic set of G contains at least one vertex from each corner path.

Proof. Without loss of generality, we assume the x -coordinates of all vertices of P are the same. By Observation 3, the set $\{v \in V(G) : v \notin V(P), N(v) \cap P \neq \emptyset\}$ induces a path Q and the x -coordinates of all vertices in Q are the same. Now consider any two vertices $a, b \in V(G) \setminus V(P)$ and with a path P' between a and b that contains one of the end-vertices, say u , of P . Observe that P' can be expressed as $P' = a c_1 c_2 \dots c_t d f_1 f_2 \dots f_{t'} u g h_1 h_2 \dots h_{t''} b$ such that $\{d, g\} \subseteq V(Q)$ and $\{f_1, f_2, \dots, f_{t'}\} \subseteq V(P)$. Then there is a path $P'' = a c_1 c_2 \dots c_t d f'_2 \dots f'_{t'} g h_1 h_2 \dots h_{t''} b$ where for $2 \leq i \leq t'$, f'_i is the vertex in Q which is adjacent to f_i in G . Observe that the length of P'' is strictly less than that of P' . Therefore $u \notin I(a, b)$ whenever $a, b \in V(G) \setminus V(P)$. Hence any geodetic set of G contains at least one vertex from P . \square

Any geodetic set of G contains all vertices of degree 1. Inspired by the above fact and Lemma 12, we define the term *corner vertex* as follows. A vertex v of G is a *corner vertex* if v has degree 1 or v is an end-vertex of some corner path. See Fig. 3(a) for an example. Observe that two corner paths may have at most one corner vertex in common. Moreover, a corner vertex cannot be in three corner paths. Therefore it follows that the cardinality of the set of corner vertices is at most $3 \cdot g(G)$.

Remark 13. Note that there are solid grid graphs whose number of corner vertices is exactly three times the geodetic number. See Fig. 3(c) for one such example.

Now we prove that the set of all corner vertices of G is indeed a geodetic set of G . We shall use the following proposition of Ekim and Erey [10].

Theorem 14 ([10]). *Let F be a graph and F_1, \dots, F_k its biconnected components. Let C be the set of cut vertices of G . If $X_i \subseteq V(F_i)$ is a minimum set such that $X_i \cup (V(F_i) \cap C)$ is a minimum geodetic set of F_i then $\cup_{i=1}^k X_i$ is a minimum geodetic set of F .*

The next observation follows from Theorem 14.

Observation 4. *Let $C(G)$ be the set of corner vertices of G and S be the set of cut vertices of G . Let $\{H_1, H_2, \dots, H_t\}$ be the set of biconnected components of G . The set $C(G)$ is a geodetic set of G if and only if $(C(G) \cap V(H_i)) \cup (S \cap V(H_i))$ is a geodetic set of H_i for all $1 \leq i \leq t$.*

From now on, $C(G)$ is the set of corner vertices of G and H_1, H_2, \dots, H_t are the biconnected components of G . Due to Theorem 14 and Observation 4, it is enough to show that for each $1 \leq i \leq t$, the set $(C(G) \cap V(H_i)) \cup (S \cap V(H_i))$ is a geodetic set of H_i . First, we introduce some more definitions below.

Let H be a biconnected component of G . Recall that each vertex of H is a pair of integers and each edge is a line segment with unit length. An edge $e \in E(H)$ is an *interior* edge if all interior points of e lie in an interior face of H . For a vertex $v \in V(H)$, let P_v denote the maximal path such that all edges of P_v are interior edges and each vertex in P_v has the same x -coordinate as v . Similarly, let P'_v denote the maximal path such that all edges of P'_v are interior edges and each vertex in P'_v has the same y -coordinate as v . A path P of H is a *red path* if (i) there exists a $v \in V(H)$ such that $P \in \{P_v, P'_v\}$ and (ii) at least one end-vertex of P is a cut-vertex or a vertex of degree 4. A vertex v of H is *red* if v lies on some red path. See Fig. 3(b) for an example.

Definition 15. *A subgraph F of H is a rectangular block if F satisfies the following properties.*

1. *For any two vertices $(a_1, b_1), (a_2, b_2)$ of F , we have that any pair (a_3, b_3) with $a_1 \leq a_3 \leq a_2$ and $b_1 \leq b_3 \leq b_2$ is a vertex of F .*
2. *Let a, a' be the maximum and minimum x -coordinates of the vertices in F . The x -coordinate of any red vertex of F must be equal to a or a' . Similarly, let b, b' be the maximum and minimum y -coordinates of the vertices in F . The y -coordinate of any red vertex of F must be equal to b or b' .*

Observe that H can be decomposed into rectangular blocks such that each non-red vertex belongs to exactly one rectangular block. See Fig. 3(b) for an example. Let B_1, B_2, \dots, B_k be a decomposition of H into rectangular blocks. Recall that $C(G)$ is the set of corner vertices of G and S is the set of cut vertices of G . We have the following lemma.

Lemma 16. *For each $1 \leq i \leq k$, there are two vertices $x_i, y_i \in (C(G) \cap V(H)) \cup (S \cap V(H))$ such that $V(B_i) \subseteq I(x_i, y_i)$.*

Proof. Let $X \in \{B_1, B_2, \dots, B_k\}$ be an arbitrary rectangular block. A vertex v of X is a *northern* vertex if the y -coordinate of v is maximum among all vertices of X . Analogously, *western* vertices, *eastern* vertices and *southern* vertices are defined. A vertex of X is a boundary vertex if it is either northern, western, southern or an eastern vertex of X . Let $nw(X)$ be the vertex of X which is both a northern vertex and a western vertex. Similarly, $ne(X)$ denotes the vertex which is both northern vertex and eastern vertex, $sw(X)$ denotes the vertex of X which is both southern and western vertex and $se(X)$ denotes the vertex of X which is both southern and eastern vertex.

First we prove the lemma assuming that all boundary vertices of X are red vertices. Let a (resp. b) denote the vertex with minimum y -coordinate such that P_a (resp. P_b) contains $sw(X)$ (resp. $se(X)$). Similarly, let c (resp. d) denote the vertex with maximum y -coordinate such that P_c (resp. P_d) contains $nw(X)$ (resp. $ne(X)$). Let a' (resp. c') denote the vertex with minimum x -coordinate such that $P_{a'}$ (resp. $P_{c'}$) contains $sw(X)$ (resp. $nw(X)$). Let b' (resp. d') denote the vertex with maximum x -coordinate such that $P_{b'}$ (resp. $P_{d'}$) contains $se(X)$ (resp. $ne(X)$). Observe that the vertices $a', a, b, b', d', d, c, c'$ lie on the exterior face of the embedding.

For two vertices $i, j \in \{a', a, b, b', d', d, c, c'\}$, let P_{ij} denote the path between i, j that can be obtained by traversing the exterior face of the embedding in the counter-clockwise direction starting from i . Observe that, if both $P_{a'a}$ and $P_{d'd}$ (resp. $P_{bb'}$ and $P_{cc'}$) contain a corner or cut vertex each, say f, f' , then $\{sw(X), ne(X)\} \subseteq I(f, f')$ (resp. $\{nw(X), se(X)\} \subseteq I(f, f')$) and therefore $V(X) \subseteq I(f, f')$. Now consider the case when at least one of the paths in $\{P_{a'a}, P_{d'd}\}$ does not contain any corner vertex or cut vertex and when at least one of the paths in $\{P_{bb'}, P_{cc'}\}$ does not contain any corner vertex or cut vertex. Due to symmetry of rotation and reflection on grids, without loss of generality we can assume that both $P_{a'a}$ and $P_{bb'}$ have no corner vertex or cut vertex. Observe that in this case there must be a corner vertex f in P_{ab} whose x -coordinate is the same as that of b and therefore of $se(X)$. If $P_{cc'}$ contains a corner vertex f' , then $\{nw(X), se(X)\} \subseteq I(f, f')$ and therefore $V(X) \subseteq I(f, f')$. Otherwise, there must be a corner vertex f' in $P_{c'a'}$ whose y -coordinate is the same as that of c' and therefore of $nw(X)$. Hence we have $\{nw(X), se(X)\} \subseteq I(f, f')$ and therefore $V(X) \subseteq I(f, f')$ in this case also.

Now we consider the case when there are some non-red boundary vertices of X . Let v be a non-red vertex of X . Without loss of generality, we can assume that v is a western vertex of X . Now we redefine the vertices $a, a', b, b', c, c', d, d'$ as follows. Let $a' = sw(X)$, $c' = nw(X)$ and a (resp. b) be the vertex with minimum y -coordinate such that there is a path from a to $sw(X)$ (resp. from b to $se(X)$) containing vertices with the same x -coordinate as that of $sw(X)$ (resp. $se(X)$). Similarly, let c (resp. d) be the vertex with maximum y -coordinate such that there is a path from c to $nw(X)$ (resp. from d to $ne(X)$) containing vertices with the same x -coordinate as that of $nw(X)$ (resp. $ne(X)$).

Finally, let d' (resp. b') be the vertex with maximum x -coordinate such that there is a path from d' to $ne(X)$ (resp. from b' to $se(X)$) containing vertices with the same y -coordinate as that of $ne(X)$ (resp. $se(X)$). Using similar arguments on the paths P_{ij} with $i, j \in \{a', a, b, b', d', d, c, c'\}$ as before, we can show that there exists corner vertices f, f' such that $V(X) \subseteq I(f, f')$. So we have the proof. \square

By Observation 4 and Lemma 16, $C(G)$ is a geodetic set of G .

Time Complexity: If the grid embedding of G is given as part of the input, then the set of corner vertices can be computed in $O(|V(G)|)$ time by simply traversing the exterior face of the embedding. Otherwise, the set of corner vertices can be computed in $O(|V(G)|)$ time as follows (we shall only describe the procedure to find corner vertices of degree two as the other case is trivial). Let H be a biconnected component of G , v be a vertex of H having degree 2 and u_0, x_0 be its neighbours. If both u_0 and x_0 have degree 4, then v is not a corner vertex. Moreover, if at least one of u_0 and x_0 have degree 2 then v is a corner vertex. Otherwise, apply the following procedure. Assume u_0 has degree 3 and denote v as u_{-1} for technical reasons. Set $i = 0$. As H is a biconnected solid grid graph, u_i and x_i must have exactly one common neighbour which is different from u_{i-1} . Denote this vertex as x_{i+1} . Let u_{i+1} be the neighbour of u_i different from both x_{i+1} and u_{i-1} . If $deg_H(u_{i+1}) = 4$ or u_{i+1} is a cut vertex in G then terminate. If $deg_G(u_{i+1}) = 2$ then v is a corner vertex. Otherwise, set $i = i + 1$ and repeat the above steps. Observe that, when the above procedure terminates either we know that v is a corner vertex or there is no corner path that contains both u_0 and v . Now swapping roles of u_0 and x_0 in the above procedure, we can decide if v is a corner vertex. We can find all the corner vertices of H by applying the above procedure to all vertices of degree 2 of H . Similarly by applying the above procedure to all the biconnected components of G , we can find all corner vertices. Notice that, the total running time of the algorithm remains linear in the number of vertices of G .

This completes the proof of Theorem 5.

4 Conclusion

In this paper, we studied the computational complexity of MGS in various graph classes. We proved that MGS remains NP-hard on planar graphs and line graphs. We also gave an $O(\sqrt[3]{n} \log n)$ -approximation algorithm for MGS on general graphs and proved that unless $P = NP$, there is no polynomial time $o(\log n)$ -approximation algorithm for MGS even on graphs with diameter 2. This motivates the following questions.

Question 1. Are there constant factor approximation algorithms for MGS on planar graphs and line graphs?

Question 2. Is there a $O(\log n)$ -approximation algorithm for MGS on general graphs ?

Acknowledgements. The authors acknowledge the financial support from the IFCAM project “Applications of graph homomorphisms” (MA/IFCAM/18/39). Florent Foucaud is supported by the ANR project HOSIGRA (ANR-17-CE40-0022). We thank Ajit Diwan for helpful discussions.

References

1. Atici, M.: Computational complexity of geodetic set. *Int. J. Comput. Math.* **79**(5), 587–591 (2002)
2. Buckley, F., Harary, F.: Geodetic games for graphs. *Quaestiones Math.* **8**(4), 321–334 (1985)
3. Bueno, L.R., Penso, L.D., Protti, F., Ramos, V.R., Rautenbach, D., Souza, U.S.: On the hardness of finding the geodetic number of a subcubic graph. *Inf. Process. Lett.* **135**, 22–27 (2018)
4. Călinescu, G., Dumitrescu, A., Pach, J.: Reconfigurations in graphs and grids. *SIAM J. Discrete Math.* **22**(1), 124–138 (2008)
5. Chlebík, M., Chlebíková, J.: Approximation hardness of dominating set problems in bounded degree graphs. *Inf. Comput.* **206**(11), 1264–1275 (2008)
6. Chuzhoy, J., Kim, D.H.K.: On approximating node-disjoint paths in grids. In: APPROX/RANDOM, pp. 187–211. Schloss Dagstuhl-Leibniz-Zentrum fuer Informatik (2015)
7. Dinur, I., Steurer, D.: Analytical approach to parallel repetition. In: STOC, pp. 624–633. ACM (2014)
8. Dourado, M.C., Protti, F., Rautenbach, D., Szwarcfiter, J.L.: Some remarks on the geodetic number of a graph. *Discrete Math.* **310**(4), 832–837 (2010)
9. Dourado, M.C., Protti, F., Szwarcfiter, J.L.: On the complexity of the geodetic and convexity numbers of a graph. In: ICDM, vol. 7, pp. 101–108. Ramanujan Mathematical Society (2008)
10. Ekim, T., Erey, A.: Block decomposition approach to compute a minimum geodetic set. *RAIRO-Oper. Res.* **48**(4), 497–507 (2014)
11. Ekim, T., Erey, A., Hegernes, P., van’t Hof, P., Meister, D.: Computing minimum geodetic sets of proper interval graphs. In: Fernández-Baca, D. (ed.) LATIN 2012. LNCS, vol. 7256, pp. 279–290. Springer, Heidelberg (2012). https://doi.org/10.1007/978-3-642-29344-3_24
12. Farber, M., Jamison, R.E.: Convexity in graphs and hypergraphs. *SIAM J. Algebraic Discrete Methods* **7**(3), 433–444 (1986)
13. Garey, M.R., Johnson, D.S.: *Computers and Intractability*, vol. 29. W.H.Freeman, New York (2002)
14. Gerber, M.U., Lozin, V.V.: Robust algorithms for the stable set problem. *Graphs Comb.* **19**(3), 347–356 (2003)
15. Gerstel, O., Zaks, S.: A new characterization of tree medians with applications to distributed sorting. *Networks* **24**(1), 23–29 (1994)
16. Gregori, A.: Unit-length embedding of binary trees on a square grid. *Inf. Process. Lett.* **31**(4), 167–173 (1989)
17. Guruswami, V.: Maximum cut on line and total graphs. *Discrete Appl. Math.* **92**(2–3), 217–221 (1999)
18. Harary, F., Loukakis, E., Tsouros, C.: The geodetic number of a graph. *Math. Comput. Modell.* **17**(11), 89–95 (1993)
19. Haynes, T.W., Henning, M., Tiller, C.A.: Geodetic achievement and avoidance games for graphs. *Quaestiones Math.* **26**(4), 389–397 (2003)

20. Itai, A., Papadimitriou, C.H., Szwarcfiter, J.L.: Hamilton paths in grid graphs. *SIAM J. Comput.* **11**(4), 676–686 (1982)
21. Mezzini, M.: Polynomial time algorithm for computing a minimum geodetic set in outerplanar graphs. *Theoret. Comput. Sci.* **745**, 63–74 (2018)
22. Mitchell, S.L.: Another characterization of the centroid of a tree. *Discrete Math.* **24**(3), 277–280 (1978)
23. Sardroud, A.A., Bagheri, A.: An approximation algorithm for the longest cycle problem in solid grid graphs. *Discrete Appl. Math.* **204**, 6–12 (2016)
24. Tirodkar, S., Vishwanathan, S.: On the approximability of the minimum rainbow subgraph problem and other related problems. *Algorithmica* **79**(3), 909–924 (2017)
25. Wu, B.Y.: A $7/6$ -approximation algorithm for the max-min connected bipartition problem on grid graphs. In: Akiyama, J., Bo, J., Kano, M., Tan, X. (eds.) *CGGA 2010*. LNCS, vol. 7033, pp. 188–194. Springer, Heidelberg (2011). https://doi.org/10.1007/978-3-642-24983-9_19
26. Yannakakis, M., Gavril, F.: Edge dominating sets in graphs. *SIAM J. Appl. Math.* **38**(3), 364–372 (1980)
27. Zhang, W., Liu, Y.: Approximating the longest paths in grid graphs. *Theoret. Comput. Sci.* **412**(39), 5340–5350 (2011)