

Centroidal Bases in Graphs

Florent Foucaud

Universitat Politècnica de Catalunya, Carrer Jordi Girona 1-3, 08034 Barcelona, Spain

Department of Mathematics, University of Johannesburg, Auckland Park 2006, South Africa

LAMSADE—CNRS UMR 7243, PSL, Université Paris-Dauphine, F-75016 Paris, France

Ralf Klasing

CNRS—LaBRI UMR 5800—Université de Bordeaux, F-33400 Talence, France

Peter J. Slater

Computer Science Department, University of Alabama in Huntsville, Huntsville, Alabama 35899

Mathematical Sciences Department, University of Alabama in Huntsville, Huntsville, Alabama 35899

We introduce the notion of a centroidal locating set of a graph G , that is, a set L of vertices such that all vertices in G are uniquely determined by their relative distances to the vertices of L . A centroidal locating set of G of minimum size is called a centroidal basis, and its size is the centroidal dimension $\text{CD}(G)$. This notion, which is related to previous concepts, gives a new way of identifying the vertices of a graph. The centroidal dimension of a graph G is lower- and upper-bounded by the metric dimension and twice the location-domination number of G , respectively. The latter two parameters are standard and well-studied notions in the field of graph identification. We show that for any graph G with n vertices and maximum degree at least 2, $(1 + o(1)) \frac{\ln n}{\ln \ln n} \leq \text{CD}(G) \leq n - 1$. We discuss the tightness of these bounds and in particular, we characterize the set of graphs reaching the upper bound. We then show that for graphs in which every pair of vertices is connected via a bounded number of paths, $\text{CD}(G) = \Omega(\sqrt{|E(G)|})$, the bound being tight for paths and cycles. We finally investigate the computational complexity of determining $\text{CD}(G)$ for an input graph G , showing that the problem is hard and cannot even be approximated efficiently up to a factor of $o(\log n)$. We also give an $O(\sqrt{n \ln n})$ -approximation algorithm. © 2014 Wiley Periodicals, Inc. NETWORKS, Vol. 000(00), 000–000 2014

Keywords: graph theory; centroidal dimension; metric dimension; location-domination; identification

Received June 2013; accepted June 2014

Correspondence to: F. Foucaud; e-mail: florent.foucaud@gmail.com

Contract grant sponsor: Programme IdEx Bordeaux—CPU; Contract grant number: ANR-10-IDEX-03-02

DOI 10.1002/net.21560

Published online in Wiley Online Library (wileyonlinelibrary.com).

© 2014 Wiley Periodicals, Inc.

1. INTRODUCTION

A large body of work has evolved concerning the problem of identifying an “intruder” vertex in a graph. As examples, one might seek to identify a malfunctioning processor in a multiprocessor network, or the location of an intruder such as a thief, saboteur, or fire in a graph-modeled facility. In this article, we introduce the new model of centroidal detection as such a graph identification problem.

An early model considered the case where one could place detection devices like sonar or Long Range Navigation (LORAN) stations at vertices in a graph; each detection device could determine the distance to the intruder’s vertex location. As introduced independently in Slater [25] and Harary and Melter [11], vertex set $L = \{w_1, \dots, w_k\} \subseteq V(G)$ is a *locating set* (also called *resolving set* in the literature) if for each vertex $v \in V(G)$, the (ordered) k -tuple $(d(v, w_1), \dots, d(v, w_k))$ of distances between the detector’s locations and the intruder vertex v uniquely determines v . A minimum cardinality locating set is called a *metric basis* (also called *reference set* in the literature), and its order is the *metric dimension* of G , denoted by $\text{MD}(G)$. Other studies involving metric bases include for example [1, 3, 12, 19, 22]. Carson [2] and, independently, Delmas et al. [4] (for the latter authors, under the name of *light r -codes*) considered the case in which each detection device at w_i can only detect an intruder at distance at most r .

In another model, the presence of any edge $\{u, v\} \in E(G)$ indicates that a detection device at u is able to detect an intruder at v . Let us denote by $N(u)$ and $N[u]$ the open and the closed neighborhood of vertex u , respectively. A set $D \subseteq V(G)$ is a *dominating set* if $\cup_{u \in D} N[u] = V(G)$.



FIG. 1. Centroidal basis $\{x_1, x_3, x_6, x_8\}$ for path P_8 .

Clearly, in the latter model, if every possible intruder location must be detectable, the set of detector locations must form a dominating set.

The concepts of locating and dominating set were merged in Slater [28, 29]: when a detection device at vertex u can distinguish between there being an intruder at u or at a vertex in $N(u)$ (but which precise vertex in $N(u)$ cannot be determined), then we have the concept of a *locating-dominating set*. More precisely, a set D of vertices is locating-dominating if it is dominating and every vertex in $V(G) \setminus D$ is dominated by a distinct subset of D . The minimum cardinality of a locating-dominating set of graph G is denoted $LD(G)$. When one can only decide if there is an intruder somewhere in $N[u]$, one is interested in an *identifying code*, as introduced by Karpovsky et al. [18]. Haynes et al. [13] added the condition that the locating-dominating set or identifying code not have any isolated vertices. When a detection device at u can determine that an intruder is in $N(u)$, but will not report if the intruder is at u itself, one is interested in *open-locating-dominating sets*, as introduced for the hypercube by Honkala et al. [15] and for all graphs by Seo and Slater [23, 24]. A bibliography of related papers is maintained by Lobstein [20].

In what follows, we will denote the path and the cycle on n vertices by P_n and C_n , respectively.

In this article, we introduce the study of *centroidal bases*. In this model, we assume that detection devices have unlimited range, as for metric bases. However, exact distances to the intruder are not known, but for detection devices u, v , if there is an intruder at vertex x , then the presence of the intruder in the graph is determined earlier by u than by v when u is closer to x than v , that is, when $d(u, x) < d(v, x)$. When $d(u, x) = d(v, x)$, we assume that u and v report simultaneously.

For example, consider $S = \{x_1, x_3, x_6, x_8\} \subseteq V(P_8)$ (see Fig. 1). For vertex x_4 , the order in which the detectors of S report is (x_3, x_6, x_1, x_8) because $d(x_4, x_3) = 1 < d(x_4, x_6) = 2 < d(x_4, x_1) = 3 < d(x_4, x_8) = 4$. For vertex x_2 , we have $d(x_2, x_1) = d(x_2, x_3)$, hence the order of reporting is $(\{x_1, x_3\}, x_6, x_8)$. The smallest size of a set $S \subseteq V(P_8)$ for which the order of reporting uniquely identifies each vertex is, in fact, four.

1.1. Medians and Centroids

In 1869, Jordan [17] showed that each of the centers and the (branch weight) centroid of a tree either consists of one vertex, or of two adjacent vertices. The *eccentricity* of vertex u is the maximum distance from u to another vertex in graph G , $e(u) = \max\{d(u, v), v \in V(G)\}$, and the *center* $\mathcal{C}(G)$ of G is the set of vertices of minimum eccentricity. For a tree T , the *branch weight* of u , $\text{bw}(u)$, is the maximum number of edges in a subtree with u as an endpoint. The *branch weight*

centroid of T is the set $\mathcal{S}_{\text{bw}}(T)$ of vertices with minimum branch weight in T .

In 1964, Hakimi [9] considered two facility location problems, one involving the center. The second one involved the *distance* $d(u) = \sum_{v \in V(G)} d(u, v)$, measuring the total response time at u (this notion was called *status* of v by Harary in 1959 [10]). The *median* of G , $\mathcal{M}(G) = \{u \in V(G), \forall v \in V(G), d(u) \leq d(v)\}$, is the set of vertices of minimum distance in G . In 1968, Zelinka [33] showed that for any tree T , $\mathcal{M}(T) = \mathcal{S}_{\text{bw}}(T)$, which seemed to imply that the median would be a good generalization of the branch weight centroid of a tree to an arbitrary graph. However (with details in Slater [27, 30] and Slater and Smart [31]), note that components of $T - u$ of the same order, one being a path and the other a star, contribute the same value of a branch weight, but have much different distances. In trying to keep closer to the spirit of what the branch weight centroid measures, the *centroid* $\mathcal{S}(G)$ of an arbitrary graph G was defined in terms of competitive facility location [26]. For facilities located at vertices u and v , a customer at vertex x is interested in which of the facilities is the closer. As defined in Slater [26], the set $V_{u,v} = \{x \in V(G), d(x, u) < d(x, v)\}$ is the set of vertex customer locations strictly closer to u than to v . Then, $f(u, v) = |V_{u,v}| - |V_{v,u}|$ rates how well u does as a facility location, in comparison to v . Letting $f(u) = \min\{f(u, v), v \in V(G) - u\}$, the *centroid* of a graph G is the set $\mathcal{S}(G) = \{u \in V(G), \forall v \in V(G) - u, f(u) \geq f(v)\}$. When G is a tree, the centroid and branch weight centroid are easily seen to coincide. Interestingly, there are graphs G for which $f(u) < 0$ for all $u \in V(G)$.

Note that in our context of centroidal bases, detectors located at u and v enable us to determine if an intruder is in $V_{u,v}$, in $V_{v,u}$, or in $V(G) - V_{u,v} - V_{v,u} = \{x \in V(G), d(x, u) = d(x, v)\}$.

1.2. Centroidal Detection

Let $B = \{w_1, \dots, w_k\} \subseteq V(G)$ be a set of vertices of graph G with detection devices located at each w_i . As noted, we will assume that each detection device has an unlimited range—an intruder entering at any vertex x will, at some point, have its presence noted at each w_i . Simply the presence will be noted, with no information about the location of the intruder. In particular, unlike in the setting of metric bases, $d(x, w_i)$ will not be known. However, the time it takes before w_i detects the intruder at x will be an increasing function of the distance $d(x, w_i)$. That is, w_i will indicate an intruder presence before w_j whenever $d(x, w_i) < d(x, w_j)$ (in our previous terminology, $x \in V_{w_i, w_j}$). We will say that x is *located* first by w_i , and then by w_j . Thus, each vertex x has a rank ordering $r(x)$ of the elements of a partition of B (in fact, $r(x)$ is an *ordered partition* of B , that is, an ordered set of disjoint subsets of B whose union is B). These ordering lists all the elements of B in nondecreasing order by their distance from x , with ties noted. Note that the number of ordered partitions of a set B of k elements is the k -th *ordered Bell number*, denoted $b(k)$ (see the book of Wilf [32, section 5.2, Example 1]).

For $B = \{x_1, x_3, x_6, x_8\}$ in path P_8 (see Fig. 1), $r(x_1) = (x_1, x_3, x_6, x_8)$, $r(x_2) = (\{x_1, x_3\}, x_6, x_8)$, $r(x_3) = (x_3, x_1, x_6, x_8)$, $r(x_4) = (x_3, x_6, x_1, x_8)$, $r(x_5) = (x_6, x_3, x_8, x_1)$, $r(x_6) = (x_6, x_8, x_3, x_1)$, $r(x_7) = (\{x_6, x_8\}, x_3, x_1)$, and $r(x_8) = (x_8, x_6, x_3, x_1)$.

Definition 1. Vertex set $B \subseteq V(G)$ is called a centroidal locating set of graph G if $r(x) \neq r(y)$ for every pair x, y of distinct vertices. A centroidal basis of G is a centroidal locating set of minimum cardinality. The centroidal dimension of G , denoted $CD(G)$, is the cardinality of a centroidal basis.

In our example, $B = \{x_1, x_3, x_6, x_8\}$ is the unique centroidal basis of path P_8 , and $CD(P_8) = 4$.

Observe that every graph G has a centroidal locating set, for example, $V(G)$: each vertex x is the only vertex having x as the first element of $r(x)$.

A useful reformulation of the definition of a centroidal locating set is as follows:

Observation 2. A set B of vertices of a graph $V(G)$ is a centroidal locating set if and only if for every pair x, y of distinct vertices of $V(G)$, there exist two vertices b_1, b_2 in B such that either $d(x, b_1) \leq d(x, b_2)$ but $d(y, b_1) > d(y, b_2)$, or $d(y, b_1) \leq d(y, b_2)$ and $d(x, b_1) > d(x, b_2)$ (in other words, $y \in V_{b_2, b_1}$ but not x , or $x \in V_{b_2, b_1}$ but not y).

1.3. Structure of the Paper

We start in section 2 by stating some preliminary observations and lemmas, and by giving bounds on parameter CD involving the order, the diameter, and other parameters of graphs; in particular, we show that $(1 + o(1)) \frac{\ln n}{\ln \ln n} \leq CD(G) \leq n - 1$ when G has n vertices and maximum degree at least 2.

In section 3, we discuss the tightness of the two aforementioned bounds by constructing graphs with small centroidal dimension, and by fully characterizing the graphs having centroidal dimension $n - 1$.

In section 4, we give a lower bound $CD(G) = \Omega\left(\sqrt{\frac{m}{k}}\right)$ when G has m edges and every pair of vertices is connected by a small number, k , of paths. We show that the bound is tight (up to a constant factor) for paths and cycles.

Finally, in section 5, we discuss the computational complexity of finding a centroidal basis; we show that for graphs with n vertices, it is NP-hard to compute an $o(\ln n)$ -approximate solution, and describe an $O(\sqrt{n \ln n})$ -approximation algorithm. We also remark that the problem is fixed-parameter-tractable when parameterized by the solution size.

2. PRELIMINARIES AND BOUNDS

In this section, we give a series of preliminary lemmas and bounds for parameter CD that will prove useful later on, and also help the reader become familiar with some of the aspects of the problem.

2.1. Preliminary Results

We state a few lemmas that will prove very useful in the study of centroidal locating sets.

Lemma 3. Let G be a graph. The following statements are true:

- (a) If u is a vertex of degree 1, then any centroidal locating set of G contains u .
- (b) If u is a vertex of degree 1 having a neighbor v of degree 2, then any centroidal locating set of G contains either v or a neighbor of v other than u .
- (c) If u, v are two vertices with $N(u) = N(v)$ or $N[u] = N[v]$, then any centroidal locating set of G contains at least one of u and v .

Proof. (a): Otherwise, u and its neighbor are not distinguished.

(b) and (c): Otherwise, u and v are not distinguished. ■

Lemma 4. Let S be a set of vertices of a graph G such that for each $u \in S$, $|N(u) \setminus S| \geq 2$ and such that for each $u, v \in S$, $N(u) \setminus S \neq N(v) \setminus S$. Then $V(G) \setminus S$ is a centroidal locating set of G .

Proof. Let $B = V(G) \setminus S$. Every vertex of B is first located by itself, while every vertex of S is first located by a distinct set of at least two vertices of B . ■

Note that in particular, Lemma 4 shows that for any vertex u of degree at least 2 in a graph G , $V(G) \setminus \{u\}$ is a centroidal locating set of G .

2.2. Bounds

We now provide some lower and upper bounds for the value of parameter CD .

Theorem 5. Let G be a graph on n vertices with maximum degree at least 2. Then

$$(1 + o(1)) \frac{\ln n}{\ln \ln n} \leq CD(G) \leq n - 1.$$

Proof. Lemma 4 immediately implies the upper bound. For the lower bound, assume that B is a centroidal basis of size $k = CD(G)$, and G has n vertices. Then, to each vertex of G , one can assign a distinct ordered partition of B . It is known that the number of ordered partitions of a set of k elements, the ordered Bell number $b(k)$, is approximated by

$$b(k) \sim \frac{k!}{2(\ln 2)^{k+1}} + O(0.16^k k!),$$

See Wilf [32, section 5.2, Example 1]. It is clear that $n \leq b(k)$. Let us assume that B is a centroidal basis of G of size k , with $n = b(k)$: for large enough n , we have $n = k!(c^{k+1})$ for some constant c . Taking the logarithm of both sides we get

$\ln n = \ln(k!) + (k + 1) \ln(c)$. Using Stirling's approximation $\ln(k!) = k \ln(k) - k + O(\ln(k))$, we obtain

$$\ln n = (1 + o(1))k \ln(k). \quad (1)$$

Hence, $k = (1 + o(1)) \frac{\ln n}{\ln(k)}$; again taking the logarithm, we get

$$\begin{aligned} \ln(k) &= \ln \ln n - \ln \ln(k) + \ln(1 + o(1)) \\ &= (1 + o(1)) \ln \ln n. \end{aligned} \quad (2)$$

Merging Equalities (1) and 2, we get

$$k = (1 + o(1)) \frac{\ln n}{\ln \ln n}.$$

Since in G , there cannot be any smaller centroidal locating set than B , the bound follows. ■

The considerations of Theorem 5 can be strengthened if we assume that the distances of a vertex to vertices in B are bounded. The following result was already known in the context of the metric dimension; see for example, Khuller et al. [19] or Chartrand et al. [3].¹

Proposition 6. *Let G be a graph with a centroidal locating set B of size k and let D be an integer. If for every vertex $u \in V(G) \setminus B$ and for every vertex $b \in B$, $d(u, b) \leq D$, then $n \leq k + D^k$ and hence $CD(G) \geq \log_D(n) - 1$. In particular, this holds if G has diameter D .*

Proof. Let $u \in V(G) \setminus B$ be a vertex. Since every vertex of B is at distance at most D from u , $r(u)$ contains at most D sets. There are exactly D^k different ordered partitions of B into at most D sets (this number is equal to the number of words of length k over an alphabet of size D), hence there can be at most D^k vertices in $V(G) \setminus B$, and hence $k + D^k$ vertices in G . ■

The bound of Proposition 6 was improved by Hernando et al. [14]:

Theorem 7 ([14]). *Let G be a graph on n vertices with diameter $D \geq 2$ and $CD(G) = k \geq 1$. Then $n \leq (\lfloor \frac{2D}{3} \rfloor + 1)^k + k \sum_{i=1}^{\lfloor D/3 \rfloor} (2i - 1)^{k-1}$.*

We improve the bound in Theorem 7 for $D \leq 3$:

Theorem 8. *Let G be a graph on n vertices with diameter $D \in \{2, 3\}$ and let $CD(G) = k$. If $D=2$ and $k \geq 1$, $n \leq 2^k + k - 1$. If $D=3$ and $k \geq 5$, $n \leq 3^k - 2^{k+1} + 2$.*

Proof. Let B be a centroidal locating set of G .

- $D=2$ and $k \geq 1$. By Theorem 7 we have $n \leq 2^k + k$. However, observe that for every vertex $v \notin B$, $r(v) = (N(v) \cap B, B \setminus N(v))$. But there can only be 2^k distinct sets $N(v) \cap B$, and moreover if $N(v) = B$ and $N(w) = \emptyset$ we have $r(v) = r(w) = (B)$. Hence $|V(G) \setminus B| \leq 2^k - 1$ and we are done.
- $D=3$ and $k \geq 5$. Since the diameter is 3, for every vertex v , $r(v)$ has at most three components, unless $v \in B$, then it may have four. Moreover, if $v \notin B$, then $r(v)$ either has one component (then $r(v) = (B)$), or two (then $r(v) = (S, B \setminus S)$ with $1 \leq |S| \leq |B| - 1$), or three (then $r(v) = (S, T, B \setminus (S \cup T))$ with $S \cap T = \emptyset$ and $1 \leq |T| \leq |B \setminus S|$).

Therefore, we have

$$|V(G) \setminus B| \leq 1 + \sum_{i=1}^{k-1} \binom{k}{i} (2^{k-i} - 1) = 3^k - 2^{k+1} + 2,$$

that is, 1 plus the number of ways of choosing a nonempty subset S of B and a nonempty subset of $B \setminus S$.

Now, if for every pair b, b' in B we have $d(b, b') \leq 2$, then no vertex v has four components in $r(v)$ and so the claimed bound holds. Now, assume that there is a pair b, b' in B with $d(b, b') = 3$. This implies that for any vertex v , $r(v)$ cannot be of the form $(\{b, b'\}, T, B \setminus (\{b, b'\} \cup T))$ with S a nonempty proper subset of $T \setminus \{b, b'\}$ since otherwise we must have $d(v, b) = d(v, b') = 1$ and hence $d(b, b') \leq 2$, a contradiction. This gives us at least $k - 2 + \binom{k-2}{2}$ forbidden triples of the form $(\{b, b'\}, T, B \setminus (\{b, b'\} \cup T))$ since T can be chosen to be one of $k - 2$ possible singletons and $\binom{k-2}{2}$ possible pairs. Since $k \geq 5$, $k - 2 + \binom{k-2}{2} \geq k$ and we are done. ■

The bounds of Theorem 8 will be proved tight in section 3.

We will now relate parameter CD to parameters MD and LD .

Lemma 9. *Let G be a graph with a locating-dominating set C such that ℓ vertices from $V(G) \setminus C$ have a unique neighbor in C . Then $CD(G) \leq LD(G) + \ell$.*

Proof. We construct a centroidal locating set C' from C by adding at most ℓ vertices to C . Note that in the setting of a centroidal locating set and considering C as a potential solution, each vertex of C is located first by itself, and then by its neighborhood within C , while each vertex v of $V(G) \setminus C$ is first located by $N(v) \cap C$. Hence, any two vertices both in C or both in $V(G) \setminus C$ are distinguished. However, if for some vertex v of $V(G) \setminus C$, $N(v) \cap C = \{c_v\}$, v and c_v might not be distinguished. In that case, it is enough to add v to C to solve this problem. This does not cause any other conflict since any superset of a centroidal locating set is also a centroidal locating set. Repeating the process ℓ times completes the proof (observing that any other vertex of C is distinguished from all other vertices). ■

Using Lemma 9, we obtain the following theorem:

¹ The result was stated in terms of the metric dimension but since any centroidal locating set is also a locating set the bound holds also for the centroidal dimension.

Theorem 10. For any graph G , $MD(G) \leq CD(G) \leq 2LD(G)$.

Proof. For the first inequality, note that any centroidal locating set B is a locating set. Indeed, if two vertices were at the same distance to each vertex of B , then they would not be distinguished by their relative distances by B , a contradiction.

The second inequality is proved by Lemma 9 by observing that for any locating-dominating set C , $\ell \leq |C|$: for each $c_v \in C$, if there were two vertices of $V(G) \setminus C$ having only c_v as a neighbor in C , they would not be distinguished by C , a contradiction. ■

For graphs of diameter 2, one gets the following improvement:

Theorem 11. Let G be a graph of diameter 2. Then $LD(G) - 1 \leq MD(G) \leq CD(G) \leq 2LD(G)$.

Proof. The last two inequalities come from Theorem 10. For the first inequality, we show that any locating set L is almost a locating-dominating set. Each vertex v of $V(G) \setminus L$ has distance 1 to all elements of its neighborhood $N_v = N(v) \cap L$ in L , and distance 2 to all vertices of $L \setminus N_v$. In other words, vertices in L only distinguish vertices they are adjacent to, from nonadjacent ones. Since L is a locating set, it follows that each vertex in $V(G) \setminus L$ has a distinct neighborhood within L . Therefore, if L is dominating, it is also locating-dominating. Otherwise, there is at most one vertex that is not dominated; adding it to set L , we get a locating-dominating set of size $|L| + 1$. ■

3. TIGHTNESS OF THE BOUNDS

In this section, we discuss the tightness of some of the bounds from section 2.2.

3.1. Graphs with Small Centroidal Dimension

For $k = 1, 2$ it is easy to construct graphs G on n vertices with $CD(G) = k$ and $n = b(k)$: for $k = 1$, $b(1) = 1$ and K_1 is the only answer; for $k = 2$, $b(2) = 3$ and both P_3, K_3 are answers.

For $k = 3$, $b(3) = 13$; the two graphs of Figure 2 have 13 vertices and centroidal dimension 3 (the black vertices form a centroidal basis).

It holds that $b(4) = 75$ [21]; a more intricate construction for this case is presented in Figure 3.

We do not know such optimal examples for $k \geq 5$ (recall that $b(k)$ grows very rapidly with k : $b(5) = 541$, $b(6) = 4683$ [21]). Note that it is not possible to directly extend our example for $k = 4$ to higher values using the same idea; indeed, every two vertices from the centroidal basis B are at distance at most 3 from each other. But in our construction, the vertices whose vector is a permutation of B have a neighbor in the basis. Hence their vector r can have length at most 4, but these $k!$ vertices need to have a vector of length k .




FIG. 2. Two graphs on $b(3) = 13$ vertices with centroidal dimension 3.

3.1.1. Bounded Diameter. We now discuss the tightness of the bounds for diameter 2 and 3 of Theorem 8.

Proposition 12. For any $k \geq 4$, there is a graph G of diameter 2 with $n = 2^k + k - 1$ vertices and $CD(G) = k$.

Proof. We construct G in the following way. We let $V(G) = B \cup S$, where B, S are disjoint sets, B has k vertices, and S has $n - k = 2^k - 1$ vertices. We make sure that in the subgraph induced by B , every vertex has a neighbor and a non-neighbor, which is possible since $k \geq 4$. The set S induces a clique and the neighborhood of every vertex of S within B is distinct and nonempty.

Since G has a vertex of degree $n - 1$ (in S) the diameter is 2. For every vertex v in B , $r(v) = (\{v\}, N(v) \cap B, B \setminus N[v])$, and by our assumption on the structure of B , these three sets are nonempty. For every vertex v in S , $r(v) = (N(v) \cap B, B \setminus N(v))$. By construction all these vectors are distinct. ■

In fact, in the construction of Proposition 12, B is a locating-dominating set; similar constructions are well known in this context, see for example Slater [29]. For diameter 3, we give a more complicated construction:

Theorem 13. For any $k \geq 4$, there is a graph G of diameter 3 with $n = 3^k - 2^{k+1} + 2$ vertices and $CD(G) = k$.

Proof. We construct G as follows (see Fig. 4 for an illustration).

- Let B be an independent set of size k , which will be the centroidal locating set of G .
- Let X be a clique containing, for every subset S of B with $2 \leq |S| \leq k - 2$, a vertex $x(S)$ that is adjacent to all vertices in S . Set X has size $\sum_{i=2}^{k-2} \binom{k}{i}$.
- Let Y be an independent set containing, for every subset S of B with $1 \leq |S| \leq k - 2$ and for every proper nonempty subset T of $B \setminus S$, a vertex $y(S, T)$ (note that $1 \leq |T| \leq k - 2$). Vertex $y(S, T)$ is adjacent to all vertices of S . Moreover, if $|T| \geq 2$, $y(S, T)$ is adjacent to $x(T)$. If $T = \{t\}$ has size 1, let T' be an arbitrary size 2-subset of B formed by t and an arbitrary vertex of S , and let $y(S, T)$ be adjacent to $x(T')$. Note that set Y has size $\sum_{i=1}^{k-2} \left(\binom{k}{i} (2^{k-i} - 2) \right)$.




FIG. 3. A graph on $b(4) = 75$ vertices with centroidal dimension 4. In the central part of the figure, the four black vertices form the centroidal basis and have their vector r of the form $(a, \{b, c, d\})$; they are at distance 3 from each other. The twelve white circle-shaped vertices have their vector of the form $(a, b, \{c, d\})$. The six gray square-shaped vertices have their vector of the form $(\{a, b\}, \{c, d\})$. The unique gray double-circled vertex has its vector of the form $(\{a, b, c, d\})$. In the outer parts of the figure, the 24 gray circle-shaped vertices have their vector of the form (a, b, c, d) . The 12 white square-shaped vertices have their vector of the form $(\{a, b\}, c, d)$. Finally, the four gray double-squared vertices have their vector of the form $(\{a, b, c\}, d)$.

- Let Z be a clique of size $k + 1$ containing, for each subset S of B with $k - 1 \leq |S| \leq k$, a vertex $z(S)$ that is adjacent to the vertices in S .

The order of G is

$$\begin{aligned}
 |B| + |X| + |Y| + |Z| &= k + \sum_{i=2}^{k-2} \binom{k}{i} + \sum_{i=1}^{k-2} \binom{k}{i} (2^{k-i} - 2) \\
 &\quad + k + 1 \\
 &= 1 + \sum_{i=1}^k \binom{k}{i} (2^{k-i} - 1) \\
 &= 3^k - 2^{k+1} + 2.
 \end{aligned}$$

Furthermore, the diameter of G is exactly 3. It is at least 3: consider some vertex $y = y(S, T)$. We have $N(y) = S$, and the vertices in T are at distance 2 from y (via vertex $x(T)$).

However, all vertices of $B \setminus (S \cup T)$ are at distance 3 from y , and since T is a proper subset of $B \setminus S$, $B \setminus (S \cup T)$ is nonempty. Conversely, the diameter is at most 3. If v is any vertex of G , v is within distance 1 of some vertex b in B , and since X and Z are cliques and b has a neighbor in both X and Z , every vertex in $X \cup Z$ is within distance 2 of b and within distance 3 of v . Moreover, every two vertices in B are at distance at most 2 apart since they all share a neighbor, $z(B)$. Hence, any vertex $y(S, T) \in Y$, since it has a neighbor in B , is at distance at most 3 from any vertex in B . Finally, any vertex in Y has a neighbor in X , which is a clique; hence any two vertices in Y have distance at most 3 from each other.

It remains to check that B is a centroidal locating set. For every $b \in B$, we have $r(b) = (\{b\}, B \setminus \{b\})$. For every vertex $x = x(S)$ in X , we have $r(x) = (S, B \setminus S)$ and $2 \leq |S| \leq k - 2$. For every vertex $z = z(S)$ in Z , $r(z) = \{S, B \setminus S\}$ if $|S| = k - 1$ and $r(z(B)) = (B)$. We have now realized all possible vectors with at most two components. Finally, for




FIG. 4. The construction of Theorem 13. Gray sets are cliques.

any vertex $y = y(S, T)$ in Y , $r(y) = (S, T, B \setminus (S \cup T))$ and by the definition of Y , none of these three components is empty. This completes the proof. ■

We leave the question of the tightness of Theorems 5 and 7 open.

Question 14. *Is the bound $CD(G) \geq (1 + o(1)) \frac{\ln n}{\ln \ln n}$ asymptotically tight, that is, can we find an infinite family of graphs G with $CD(G) = O\left(\frac{\ln n}{\ln \ln n}\right)$?*

Observe that, by Proposition 6, $CD(G) \geq \log_D(n) - 1$ when G has diameter D and n vertices. Hence, to construct a graph with a centroidal locating set of size $O\left(\frac{\ln n}{f(n)}\right)$ for some $f(n) = O(\ln \ln n)$, G should have diameter $\Omega(e^{f(n)})$; in particular, for $f(n) = \ln \ln n$, the diameter should be $\Omega(\ln n)$.

It was proved by Sebő and Tannier [22] that for the hypercube Q_k with $n = 2^k$ vertices, $MD(Q_k) = \frac{2 \log_2 n}{\log_2 \log_2 n} (1 + o(1))$. In this regard, it would be interesting to determine whether the family of hypercubes is a positive answer to Question 14; determining $CD(Q_k)$ would be of independent interest.

Question 15. *What is the maximum order of a graph with centroidal dimension k and diameter $D \geq 4$?*

3.2. Graphs with Large Centroidal Dimension

A direct consequence of Theorem 5 is that a graph has centroidal dimension equal to its order if and only if it has maximum degree at most 1. We now fully characterize the set of graph with centroidal dimension of value the order minus one.

For some $n \geq 1$, K_n denotes the complete graph on n vertices. For some $a, b \geq 1$, $K_{a,b}$ denotes the complete bipartite

graph with parts of sizes a and b . Let $n \geq 4$. We denote by S_n , the graph obtained by joining K_2 to an independent set of $n-2$ vertices. We call T_n the tree obtained from P_3 by attaching $n-3$ degree 1-vertices to one of the ends of P_3 . Finally, we call U_n the graph obtained from K_3 by attaching $n-3$ degree 1-vertices to one of the vertices of K_3 .

In particular, $K_{1,n-1}$ is a star, $K_{2,2}$ is the cycle C_4 , S_4 is the diamond graph, and T_4 is the path P_4 . See Figure 5 for illustrations of these graphs (black vertices belong to a centroidal basis).

Proposition 16. *Let G be a graph on $n \geq 3$ vertices belonging to $\{K_n, K_{1,n-1}, K_{2,n-2}, S_n, T_n, U_n\}$. Then $CD(G) = n - 1$.*

Proof. By Lemma 4 applied on a single degree 2-vertex, it is easily seen that in each case, $CD(G) \leq n - 1$. If G is isomorphic to K_n , the lower bound is directly implied by Lemma 3(c).

If G is isomorphic to $K_{1,n-1}$, then by Lemma 3(a) any centroidal locating set contains all $n-1$ leaves of $K_{1,n-1}$.

If G is isomorphic to $K_{2,n-2}$ or to S_n , by Lemma 3(c) at least $n-3$ vertices of degree 2 belong to any centroidal locating set, as well as one of the other two vertices. However, if no further vertex does belong to the centroidal locating set, then the vertex of degree 2 that is not in the set is not distinguished from its neighbor in the set, a contradiction.

If G is isomorphic to T_n , by Lemma 3(a) all $n-2$ vertices of degree 1 belong to any centroidal locating set. Moreover, one further vertex also does according to Lemma 3(b).

Finally, if G is isomorphic to U_n , again by Lemma 3(a) all $n-3$ vertices of degree 1 belong to any centroidal locating set, as well as one degree 2-vertex of the triangle by Lemma 3(c). If no further vertex does belong to a centroidal locating set, then the two degree 2-vertices of the triangle are not distinguished, a contradiction. ■

In fact, the graphs from Figure 5 are the only extremal graphs, as shown in the following theorem:

Theorem 17. *Let G be a connected graph on $n \geq 3$ vertices with $CD(G) = n - 1$. Then G belongs to $\{K_n, K_{1,n-1}, K_{2,n-2}, S_n, T_n, U_n\}$.*

Proof. We assume by contradiction that $CD(G) = n - 1$ but G does not belong to the list. Hence, $n \geq 5$ since all connected graphs on three or four vertices are in the list. Let u be a vertex of G of degree at least 2. By Lemma 4, $V(G) \setminus \{u\}$ is a centroidal locating set. First of all, observe that there is no vertex at distance 3 of u : for contradiction, assume that z is such a vertex and let t be a neighbor of z with $\deg(t) \geq 2$ that lies on a path from u to z . By Lemma 4, $B = V(G) \setminus \{u, t\}$ is a centroidal locating set, a contradiction.

We now assume that $\deg(u) = 2$. Let v, w be the two neighbors of u , and let x, y be two other vertices in G (they exist since $n \geq 5$). Since there is no vertex at distance 3 of u , both x, y are neighbors of at least one of v, w . First, assume




FIG. 5. The list of graphs on n vertices with centroidal dimension $n-1$.

that x is a neighbor of both v and w and that y is only a neighbor of v . Then $B = V(G) \setminus \{v, w\}$ is a centroidal locating set, a contradiction. Indeed, each vertex of B is first located by itself, then by its neighbors in B , while v, w are first located by a set of at least two vertices. Moreover, v is (in particular) first located by y , while w is not.

Hence, either all vertices other than u, v, w are adjacent to both v, w , or none is. In the first case, if all common neighbors of v, w form an independent set, G is isomorphic to either $K_{2,n-2}$ or S_n , a contradiction. Hence, there is an edge between two common neighbors of v, w , say between x, y . Then, $B = V(G) \setminus \{u, x\}$ is a centroidal locating set, a contradiction. Indeed, each vertex of B is first located by itself, then by its neighbors in B , while u, x are first located by a set of at least two vertices. Moreover, x is (in particular) first located by y , while u is not.

We now have that each vertex other than u, v, w is adjacent to exactly one of v, w . Let S_v be the set of neighbors of v other than u, w and let S_w be the set of neighbors of w other than u, v . If both $|S_v|, |S_w| \geq 1$, by Lemma 4, $V(G) \setminus \{v, w\}$ is a centroidal locating set, a contradiction. Otherwise, either G is isomorphic to T_n if v, w are nonadjacent, or to U_n otherwise, a contradiction in both cases.

By the previous discussion, G has no degree 2-vertices. Hence $\deg(u) \geq 3$. Suppose, moreover, that $N(u)$ is an independent set. Since G is not isomorphic to $K_{1,n-1}$, there are vertices at distance 2 from u . Hence u has a neighbor, v , with two such vertices as neighbors (since $\deg(v) \neq 2$): let x, y be these vertices. Then, $B = V(G) \setminus \{u, v\}$ is a centroidal locating set, a contradiction. Indeed, all vertices but u, v are located first by themselves only, whereas u is located first by $N(u) \setminus \{v\}$ and v by a set containing both x, y .

Now, we assume that $N(u)$ is a clique. Since G is not a complete graph, u has a neighbor v , having a neighbor x with $d(u, x) = 2$. Then, by Lemma 4, $V(G) \setminus \{u, v\}$ is a centroidal locating set of G , a contradiction.

Hence, $N(u)$ is neither an independent set, nor a clique: there is a vertex v in $N(u)$ with a neighbor w and a non-neighbor x , both being in $N(u)$. Since no vertex in G has degree 2, v has an additional neighbor, y . But then by Lemma 4, $V(G) \setminus \{u, v\}$ is a centroidal locating set of G , a contradiction. ■

4. GRAPHS WITH FEW PATHS CONNECTING EACH PAIR OF VERTICES

We now study parameter CD for graphs in which every pair of vertices is connected via a bounded number k of paths. We show that such graphs have centroidal dimension $\Omega\left(\sqrt{\frac{m}{k}}\right)$, where m is the number of edges. In particular, this applies to paths and cycles; for these graphs, we show that the lower bound is asymptotically tight. These cases are particularly interesting for the following reason: the metric dimension of a path or a cycle with n vertices is easily seen to be constant (1 for any path, 2 for any cycle), whereas the location-domination number is linear (roughly $\frac{2}{5}$ th of the vertices [29]). In contrast, the centroidal dimension is about the square-root of the order.

But first, the following technical lemma will be useful when showing our lower bound.

Lemma 18. *Let G be a graph with u, v two adjacent vertices, and let B be a centroidal locating set. Then, there are*

two vertices b_1, b_2 of B and a path $P : b_1 - u - v - b_2$ such that at least one of the following properties hold:

1. $\{u, v\} = \{b_1, b_2\}$ and P is the edge $\{u, v\}$;
2. $d(u, b_1) = d(u, b_2)$, $d(v, b_1) \neq d(v, b_2)$ (or, symmetrically, $d(v, b_1) = d(v, b_2)$, $d(u, b_1) \neq d(u, b_2)$) and P contains a shortest path from u to b_1 and a shortest path from v to b_2 ;
3. $d(u, b_1) + 1 = d(u, b_2)$, $d(v, b_1) = d(v, b_2) + 1$, P contains a shortest path from u to b_1 and a shortest path from v to b_2 , P has odd length and $\{u, v\}$ is the middle edge of P .

Proof. If both u, v belong to B , we are in the first case and we are done.

Otherwise, since B is a centroidal locating set, without loss of generality there are two vertices b, b' of B such that $d(u, b) \leq d(u, b')$ and $d(v, b) > d(v, b')$, or vice versa.

CASE a $d(u, b) = d(u, b')$. We show that the second case of the statement holds. If v lies on a shortest path P_u from u to one of b, b' (say, b), we are done: then no shortest path from u to b' can go through v (otherwise $d(v, b) = d(v, b')$). Hence, setting $b = b_2$ and $b' = b_1$, the concatenation of P_u and any shortest path from u to b' is a path P satisfying the desired properties.

Hence, we assume that v does not lie on a shortest path from u to b . Since $d(v, b) \neq d(v, b')$, we can assume without loss of generality that $d(v, b') \neq d(u, b') + 1$ (moreover, since $d(v, b') \leq d(u, b') + 1$, we have $d(v, b') \leq d(u, b')$). Hence, a shortest path P_v from v to b' does not go through u . Let P_u be a shortest path from b to u . If the concatenation $P_u - uv - P_v$ is a path, we are done by setting $P = P_u - uv - P_v$, $b = b_1$ and $b' = b_2$. Therefore, assume that it is not a path, and let w be the vertex closest to u appearing in both P_u and P_v : $P_u - uv - P_v$ contains a cycle going through u, v, w . Now, since no shortest path from v to b' goes through u , we have $d(v, w) \leq d(u, w)$. Also, since we assumed that v does not lie on a shortest path from u to b , in particular $d(u, w) \leq d(v, w)$. Therefore, $d(u, w) = d(v, w)$. This implies that $d(v, b') = d(u, b')$ (indeed, we had $d(v, b') \leq d(u, b')$ and now $d(u, b') \leq d(u, w) + d(w, b') = d(v, w) + d(w, b') = d(v, b')$). Hence, $d(v, b) < d(v, b')$. Let P'_u be the path obtained from the concatenation of a shortest path from u to w and the subpath of P_u from w to b' , and let P'_v be a shortest path from v to b . Then, P'_v does not contain any vertex of P'_u (indeed, if there was such a vertex t , then $d(u, b) \leq d(v, b) < d(v, b') = d(u, b')$, a contradiction). Hence, the concatenation $P = P'_v - uv - P'_u$ is a path that has the desired properties.

CASE b $d(u, b) < d(u, b')$. We show that the third case of the statement holds. Since $d(v, b) > d(v, b')$ and u, v are adjacent, $d(v, b) = d(u, b) + 1$ and $d(v, b') = d(u, b') - 1$. Hence $d(v, b') < d(v, b) < d(v, b') + 2$ and $d(u, b) < d(u, b') < d(u, b) + 2$, and $d(u, b) = d(v, b')$. We set $b_1 = b$ and $b_2 = b'$. Observe that the concatenation P of a shortest path from b_1 to u , the edge uv , and a shortest path from v to b_2 is a path from b_1 to b_2 (if it were not a path, we would

have $d(u, b) = d(u, b') = d(v, b) = d(v, b')$). Since P has the desired properties, this completes the proof. ■

In what follows, for a pair $\{u, v\}$ of vertices in a graph G , we let $k_{\text{odd}}(u, v)$ and $k_{\text{ev}}(u, v)$ be the number of odd and even (not necessarily disjoint) paths connecting u to v , respectively.

Theorem 19. Let G be a graph on n vertices and m edges such that for every pair $\{u, v\}$ of vertices, $2k_{\text{ev}}(u, v) + k_{\text{odd}}(u, v) \leq k$ for some integer k . Then, $CD(G) > \sqrt{\frac{2m}{k}}$. In particular, for every tree T , $CD(T) > \sqrt{n-1}$.

Proof. Let B be a centroidal locating set of G . To each pair u, v of adjacent vertices in G , we assign a triple $(b_1, b_2, P) = T(u, v)$ of two vertices b_1, b_2 of B and a path $P : b_1 - u - v - b_2$ satisfying one of the three properties described in Lemma 18. The assignment is done as follows:

1. if both u, v belong to B , set $T(u, v) = (u, v, uv)$ (we say that the pair u, v is of type 1);
2. otherwise, if there is a pair $\{b_1, b_2\}$ of B and a path P for which the second property of Lemma 18 holds, then set $T(u, v) = (b_1, b_2, P)$ (we say that the pair u, v is of type 2);
3. otherwise, let $T(u, v)$ consist of an arbitrary pair b_1, b_2 of vertices of B with a path P such that the third property of Lemma 18 holds (we say that the pair u, v is of type 3).

Now, for a given pair b_1, b_2 of B and a path P from b_1 to b_2 , we will upper bound the number of pairs of adjacent vertices u, v such that $T(u, v) = (b_1, b_2, P)$.

Assume first that P has odd length. If P is the edge b_1, b_2 , there is only the pair $\{u, v\} = \{b_1, b_2\}$ of type 1 with $T(u, v) = (b_1, b_2, P)$. Otherwise, assume $\{u, v\}$ is a pair of type 2 with $T(u, v) = (b_1, b_2, P)$. Then by Lemma 18, $d(u, b_1) = d(u, b_2) = \ell_1$ and the subpath of P from u to b_1 has length ℓ_1 . Since the subpath from v to b_2 is also a shortest path and $d(u, b_2) = \ell_1$, it has length ℓ_2 with $\ell_1 - 1 \leq \ell_2 \leq \ell_1 + 1$. The length of P is $\ell_1 + \ell_2 + 1$. Since it is an odd number, $\ell_2 = \ell_1$ and $\{u, v\}$ is the middle edge of P . If $\{u, v\}$ is of type 3, by Lemma 18, $\{u, v\}$ is also the middle edge of P . Hence in total there is at most one pair u, v with $T(u, v) = (b_1, b_2, P)$.

Now, assume that P has even length. Let $\{u, v\}$ be a pair with $T(u, v) = (b_1, b_2, P)$. Then, by Lemma 18, u, v cannot be of type 1 or type 3, so it must be of type 2. By the same arguments as in the previous paragraph, the length of the subpath of P from b_1 to u is ℓ_1 , and the length of the subpath of P from v to b_2 is either $\ell_1 - 1$ or $\ell_1 + 1$. In both cases, one of u, v is the middle vertex of P . Hence there can be at most two such pairs.

To summarize, we proved that for each pair b_1, b_2 of B , the number of pairs u, v with $T(u, v) = (b_1, b_2, P)$ for some path P is at most

$$2k_{\text{ev}}(b_1, b_2) + k_{\text{odd}}(b_1, b_2) \leq k.$$




FIG. 6. Illustration of the sets defined in the proof of Theorem 20.

Since in total, there are m pairs of adjacent vertices in G and each such pair is associated to exactly one of the $\binom{|B|}{2}$ pairs of B , we have $m \leq k \binom{|B|}{2} < \frac{k|B|^2}{2}$, and the bound follows.

When G is a tree, there are $n-1$ edges, and there is a unique path between any pair of vertices, hence $k \leq 2$. ■

We now show that the bound of Theorem 19 is tight up to a constant factor for paths and cycles.

Theorem 20. *Let $n \geq 3$.*

1. *If n is even, $\frac{\sqrt{2}}{2}\sqrt{n} < CD(C_n)$. If n is odd, $\frac{\sqrt{6}}{3}\sqrt{n} < CD(C_n)$. In both cases, $CD(C_n) < \frac{7\sqrt{2n}}{2} + 1$. If $n = 2\ell^2$ for some integer ℓ , then $CD(C_n) \leq \sqrt{2n} - 2$.*
2. *$\sqrt{n-1} < CD(P_n) < 6\sqrt{n-1} + 3$. If $n = (2\ell)^2 + 1$ for some integer ℓ , then $CD(P_n) \leq 2\sqrt{n-1} - 3$.*

Proof. Lower bounds. The bounds follow from Theorem 19 since paths are trees, and by observing that cycles have n edges. For even cycles, for any pair $\{u, v\}$ of vertices, $k_{\text{odd}}(u, v) \leq 2$ and $k_{\text{ev}}(u, v) \leq 2$ and $k_{\text{odd}}(u, v) + k_{\text{ev}}(u, v) = 2$, hence $2k_{\text{ev}}(u, v) + k_{\text{odd}}(u, v) \leq 4$. For odd cycles, $k_{\text{odd}}(u, v) = 1$ and $k_{\text{ev}}(u, v) = 1$, hence $2k_{\text{ev}}(u, v) + k_{\text{odd}}(u, v) \leq 3$.

Upper bounds for cycles. We first prove that for any $p, q \geq 2$, if $n = p(2q+2)$, then

$$CD(C_n) \leq p + q - 1. \quad (3)$$

Assuming that $n = 2\ell^2$, and setting $p = \ell$ and $q = \ell - 1$, Inequality (3) yields the claimed bound $CD(C_n) \leq 2\ell - 2 = \sqrt{2n} - 2$.

Let $\{x_0, \dots, x_{n-1}\}$ be the vertex set of C_n . Let us divide C_n into p portions of $2q+2$ consecutive vertices each: for $0 \leq i \leq p-1$, $R_i = \{x_{i(2q+2)}, x_{i(2q+2)+1}, \dots, x_{(i+1)(2q+2)-1}\}$. For each i , we further define two subsets of R_i as follows: $S_i = \{s_i^1, \dots, s_i^q\}$ and $T_i = \{t_i^1, \dots, t_i^q\}$, where for $1 \leq j \leq q$, $s_i^j = x_{i(2q+2)+j}$ and $t_i^j = x_{i(2q+2)+q+1+j}$. In other words, S_i contains q vertices from R_i , starting from the second one, and

T_i contains the q last vertices of R_i . Observe that the first and the $(q+2)$ -nd vertices from R_i neither belong to S_i , nor T_i .

We now define a set $B = B_0 \cup B_1$, which we claim, will be our centroidal locating set. We let $B_0 = \{b_0^0, \dots, b_0^{q-1}\}$, where for $0 \leq i \leq q-1$, $b_0^i = x_{2(i+1)}$ (that is, B_0 contains each second vertex of R_0). We let $B_1 = \{b_1^0, \dots, b_1^{p-1}\}$, where for $0 \leq i \leq p-1$, $b_1^i = x_{i(2q+2)}$ (that is, B_1 contains the first vertex of each set R_i).

An illustration of sets S_i, T_i, B_0, B_1 is given in Figure 6.

Observe that B has $p+q-1$ elements. It remains to show that B is a centroidal locating set.

First of all, notice that for any $0 \leq i \leq p-1$, $b_1^i \in B_1$ is the unique vertex that is first located by itself, and later by $\{b_1^{(i-1) \bmod p}, b_1^{(i+1) \bmod p}\}$ at the same time. For $i \neq 0$, the $(q+2)$ -nd vertex of R_i , $x_{i(2q+2)+q+1}$, is the only one that is located first by $\{b_1^i, b_1^{(i+1) \bmod p}\}$ at the same time. Similarly, x_{q+1} is the unique vertex first located by the vertices of B_0 (in some order), and then by $\{b_1^0, b_1^1\}$ at the same time. If $x \in S_0 \cup T_0$, if $x \in B_0$, x is the only vertex located first by itself only; if $x \notin B_0$, x is the only vertex located first by its two neighbors. Hence, all the previously considered vertices are distinguished from any other vertex in C_n .

Now, let u, v be a pair of vertices not yet proved to be distinguished. If $u \in S_i, v \in T_i$ ($1 \leq i \leq p-1$), then $d(u, b_1^i) < d(u, b_1^{(i+1) \bmod p})$ but $d(v, b_1^i) > d(v, b_1^{(i+1) \bmod p})$. If $u \in S_i, v \in S_{i'}$ ($1 \leq i, i' \leq p-1$), then $d(u, b_1^i) < d(u, b_1^{i'})$ but $d(v, b_1^i) > d(v, b_1^{i'})$. The case where $u \in T_i, v \in T_{i'}$ is symmetric. If $u \in S_i, v \in T_{i'}$ ($1 \leq i, i' \leq p-1$), if $i = (i'+1) \bmod p$, then $d(u, b_1^{(i+1) \bmod p}) < d(u, b_1^{i'})$ while $d(v, b_1^{(i+1) \bmod p}) > d(v, b_1^{i'})$. Otherwise, $d(u, b_1^i) < d(u, b_1^{(i'+1) \bmod p})$ while $d(v, b_1^i) > d(v, b_1^{(i'+1) \bmod p})$.

It remains to prove that for $1 \leq i \leq p-1$, any two vertices from S_i are distinguished (the case of two vertices of T_i would follow by symmetry). To see this, let $s_i^j \in S_i$. If $i \leq \left\lceil \frac{p-1}{2} \right\rceil$, $d(s_i^j, b_1^{2i}) = d(s_i^j, b_1^0) = (2q+2)i - j$, and no other vertex of S_i has this property. Similarly, if $i > \left\lceil \frac{p-1}{2} \right\rceil$, $d(s_i^j, b_1^{2i-p}) = d(s_i^j, b_1^0) = (2q+2)(p-i) + j$. This completes the proof of validity of B .

To prove the bound for all cycles, if n is not of the form $2\ell^2$, let $m = 2\ell^2$ be the integer of this form that is closest to n and such that $m \leq n: n = m + k$ for some k , and $n < 2(\ell + 1)^2$. A construction similar to the previous one can be done. Letting $p = \ell$, $q = \ell - 1$, we construct $B_1 = \{b_1^0, \dots, b_1^{p-1}\}$ with $b_1^i = x_{i(2q+2)}$ as previously; however this time, B_1 does not include any vertex from $\{x_m, \dots, x_{n-1}\}$. Instead, we let $B_0 = \{x_1, \dots, x_{2q+1}\} \cup \{x_m, \dots, x_{n-1}\}$: B_0 contains the first $2q+2$ vertices (except x_0), together with the k last vertices. It is clear that the construction works the same way as in the previous proof—we omit the details as a formal proof would be tedious.² In total, $B = B_0 \cup B_1$ has size at most $p + 2q + 1 + k = 3\ell - 1 + k$. Since $2(\ell + 1)^2 - 2\ell^2 = 4\ell + 2$, $k < 4\ell + 2$ and $|B| \leq 7\ell + 1$. Furthermore, $2\ell^2 < n$ and $\ell < \frac{\sqrt{2n}}{2}$. Hence $|B| < \frac{7\sqrt{2n}}{2} + 1$, proving the bound.

Upper bounds for paths. A similar construction as the one for cycles can be done for the case of paths: for any $p, q \geq 2$, if $n = p(2q + 2) + 1$, then $\text{CD}(P_n) \leq p + 2q - 1$. The idea of the construction is again to divide the vertex set $\{x_0, \dots, x_{n-1}\}$ into p portions of size $2q + 2$ each (vertex x_{n-1} does not belong to any such portion). For $0 \leq i \leq p$, $b_1^i = x_{i(2q+2)}$, and B_1 contains all vertices of the form b_1^i . B_0 is defined in the same way as for our construction for cycles, but now we also consider a set B'_0 similar to B_0 , but on the other end of the path. We have $|B_0 \cup B'_0 \cup B_1| = p + 1 + 2(q - 1) = p + 2q - 1$, and similar arguments as for the construction for cycles show that $B_0 \cup B'_0 \cup B_1$ is a centroidal locating set.

Hence, assuming $n = (2\ell)^2 + 1$ for some $\ell \geq 2$ and setting $p = 2\ell$ and $q = \ell - 1$, we get $\text{CD}(P_n) \leq 4\ell - 3 = 2\sqrt{n-1} - 3$.

For the general bound, once again we do not optimize the constant. Assume that n is not of the form $(2\ell)^2 + 1$, and let $m = (2\ell)^2 + 1$ be the integer of this form that is closest to n and $m \leq n$: we have $n = m + k$ for some k , and $n < (2(\ell + 1))^2 + 1$. Let $p = 2\ell$ and $q = \ell - 1$. Now, B_1 is selected as before among the first m vertices; B_1 has p elements. B_0 contains the first $2q + 2$ vertices (except x_0), and B'_0 contains the last $k + 2q + 3$ vertices (except x_m). In total, $B = B_0 \cup B'_0 \cup B_1$ has $p + (2q + 1) + (k + 2q + 2) = p + 4q + 3 + k = 6\ell - 1 + k$ vertices. Since $(2(\ell + 1))^2 + 1 - (2\ell)^2 - 1 = 6\ell + 4$, $k < 6\ell + 4$. This implies $|B| < 12\ell + 3$. Since $n > (2\ell)^2 + 1$, $\ell < \frac{\sqrt{n-1}}{2}$; hence, $|B| < 6\sqrt{n-1} + 3$. ■

5. COMPLEXITY RESULTS

Let us now turn our attention to the computational complexity of finding a small centroidal locating set, that is, the computational complexity of the following problem:

CENTROIDAL DIMENSION
 INSTANCE: A graph G .
 TASK: Find a centroidal basis of G .

² In fact we have not taken care of optimizing the construction, as here all the vertices of B_0 are probably not needed.

We have seen in Theorem 11 that for any graph of diameter 2, $\text{LD}(G) - 1 \leq \text{CD}(G) \leq 2\text{LD}(G)$. We get the following corollary, showing that CENTROIDAL DIMENSION is computationally very hard, even from the approximation point of view (recall that α -approximation algorithm for problem P is a polynomial-time algorithm for P which always outputs a solution of size no greater than α times the size of an optimal solution).

Corollary 21. CENTROIDAL DIMENSION is NP-hard to approximate within any factor $o(\ln n)$ for graphs on n vertices (even for graphs with a vertex adjacent to all other vertices, and hence diameter 2-graphs). For graphs of diameter 2, it has an $O(\ln n)$ -approximation algorithm.

Proof. Since $\text{LD}(G) - 1 \leq \text{CD}(G) \leq 2\text{LD}(G)$ and the bounds are constructive, any α -approximation algorithm ($\alpha \geq 1$) for MINIMUM LOCATING-DOMINATING SET can be transformed into an approximation algorithm of factor $2\alpha(1 + \frac{1}{\text{OPT}}) = 2\alpha(1 + o(1))$ for CENTROIDAL DIMENSION for graphs of diameter 2, and vice versa. Indeed, given an α -approximate locating-dominating set D of G , we construct a centroidal locating set of G of size at most $2|D|$. We have $2|D| \leq 2\alpha\text{LD}(G) \leq 2\alpha(\text{CD}(G) + 1) = (2\alpha + \frac{2\alpha}{\text{CD}(G)})\text{CD}(G)$. For the converse, the reasoning is similar.

This also implies that if MINIMUM LOCATING-DOMINATING SET is NP-hard to α -approximate for graphs of diameter 2 for some $\alpha \geq 2$, then CENTROIDAL DIMENSION is NP-hard to approximate within factor $(\frac{\alpha}{2} \frac{\text{OPT}}{\text{OPT}+1}) = \frac{\alpha(1-o(1))}{2}$ for graphs of diameter 2.

The positive approximation bound follows, as MINIMUM LOCATING-DOMINATING SET is well known to be $O(\ln n)$ -approximable, see for example Gravier et al. [8].

Moreover, it follows from a reduction for MINIMUM IDENTIFYING CODE in the first author's thesis [5, section 6.4] and a lemma from Gravier et al. [8] (see also Foucaud [6, 7]) that MINIMUM LOCATING-DOMINATING SET is NP-hard to approximate within a factor of $o(\ln n)$ for graphs having a vertex adjacent to all other vertices. This proves the nonapproximability bound. ■

Note that Corollary 21 fully determines the computational complexity of CENTROIDAL DIMENSION in graphs of diameter 2 from the approximation point of view. It was recently proved by Hartung and Nichterlein that the related problem METRIC DIMENSION remains NP-hard to approximate within a factor of $o(\ln n)$ even for subcubic graphs [12]. In general, it would be interesting to extend the result of Corollary 21 to other families of graphs.

As is often the case with domination or identification problems in graphs, a good way of reformulating our problem is to represent it as an instance of MINIMUM SET COVER, which is well known to be $(\ln n + 1)$ -approximable (see Johnson [16]):

MINIMUM SET COVER
 INSTANCE: A hypergraph $H = (V, E)$.

TASK: Find a minimum-size subset $C \subseteq E$ such that $\cup_{X \in C} X = V$.

For example, METRIC DIMENSION for a graph G can be expressed in this way by constructing a hypergraph $H_{MD}(G)$ on vertex set $\binom{V(G)}{2}$ with a hyperedge E_v for each vertex $v \in V(G)$ containing all pairs $\{x, y\}$ of vertices with $d(x, v) \neq d(y, v)$. Then $H_{MD}(G)$ has a set cover of size k if and only if G has a locating set of size k , as shown by Khuller et al. [19]. Hence METRIC DIMENSION is $O(\ln n)$ -approximable.

Next, we give a similar reduction for CENTROIDAL DIMENSION, but with a weaker approximation ratio.

Theorem 22. CENTROIDAL DIMENSION IS $O(\sqrt{n \ln n})$ -approximable for graphs on n vertices.

Proof. Using Observation 2, finding a centroidal locating set is equivalent to finding a set of *pairs* of vertices which identifies each pair of vertices in G —where a pair b_1, b_2 identifies x, y if $d(x, b_1) \leq d(x, b_2)$ but $d(y, b_1) > d(y, b_2)$, or $d(y, b_1) \leq d(y, b_2)$ and $d(x, b_1) > d(x, b_2)$.

Let G be a graph on n vertices. We define the hypergraph $H = H_{CD}(G)$ on vertex set $V(H) = \binom{V(G)}{2}$. For each pair $\{x, y\}$ of vertices of G , H has a hyperedge $E_{x,y}$ that contains all pairs that are identified by $\{x, y\}$.

Let C be a set cover of H . Using our previous observation, one can construct a centroidal locating set $B(C)$ of G by taking the union of all elements in the pairs that correspond to hyperedges in C : $B(C) = \{x \in V(G), \exists y \in V(G), E_{x,y} \in C\}$. Indeed, every pair of vertices of G is identified by a pair corresponding to a hyperedge of H . Hence we have

$$|B(C)| \leq 2|C|. \quad (4)$$

For the other direction, given a centroidal basis B of G , one can construct a set cover of H consisting of all $\binom{|B|}{2}$ pairs of vertices of B : each pair u, v in $V(G)$ is identified by some pair x, y in B , hence vertex $\{u, v\}$ in H is covered by the corresponding hyperedge $E_{x,y}$. Denoting by $SC(H)$ the size of an optimal set cover of H , this implies

$$SC(H) \leq \binom{|B|}{2} \leq \binom{CD(G)}{2}. \quad (5)$$

Now, to approximate CENTROIDAL DIMENSION, we construct $H = H_{CD}(G)$ from G , and apply the standard approximation algorithm for MINIMUM SET COVER [16] to get a set cover C of H of size $O(\ln n \cdot SC(H))$. By Inequalities (4) and (5), we get a centroidal locating set $B(C)$ of size at most $2|C| = O(\ln n \cdot SC(H)) = O(\ln n \cdot CD(G)^2)$.

Now, if $CD(G) \geq \sqrt{\frac{n}{\ln n}}$, we have a trivial $O(\sqrt{n \ln n})$ -approximation by selecting all vertices as a solution, since $n = O(\sqrt{n \ln n} \cdot CD(G))$.

If $CD(G) \leq \sqrt{\frac{n}{\ln n}}$, $|B(C)| = O\left(\ln n \cdot CD(G) \cdot \sqrt{\frac{n}{\ln n}}\right) = O\left(\sqrt{n \ln n} \cdot CD(G)\right)$. ■

We note that the quadratic dependence between $SC(H)$ and $CD(G)$ is necessary for Inequality (5) in the reduction of Theorem 22. Indeed, when we considered the case of cycles

in section 4, we had n special pairs to distinguish using other pairs b_1, b_2 from B , but any pair b_1, b_2 could only distinguish a small (constant) number of these n pairs. However, we could build a centroidal locating set B of size $O(\sqrt{n})$, meaning that a large fraction of the pairs from B were indeed necessary to distinguish the pairs. Hence this would lead to a set cover of H of size $\Omega(|B|^2)$ in our reduction. This suggests that one cannot improve the approximation ratio from Theorem 22 for CENTROIDAL DIMENSION using our reduction. Hence we ask the following question:

Question 23. What is the exact approximation complexity of CENTROIDAL DIMENSION?

We close the section with a remark on the parameterized complexity of CENTROIDAL DIMENSION: this problem is fixed-parameter-tractable with parameter k , the size of the solution, that is, it admits an algorithm of running time $f(k)n^{O(1)}$ for some computable function f :

Proposition 24. CENTROIDAL DIMENSION is fixed-parameter-tractable when parameterized by the size of the solution.

Proof. We know that the order n of a graph with a centroidal locating set of size k is at most $b(k)$, hence, if the input has more than $b(k)$ vertices, we answer NO. Otherwise, we enumerate all $\binom{n}{k}$ subsets of vertices of size k to check if one of them is a centroidal locating set (checking whether a given set is centroidal locating can be done in time $n^{O(1)}$). This algorithm has running time $\binom{n}{k} n^{O(1)} \leq n^{k+O(1)} = b(k)^{k+O(1)}$, which is computable and only depends on k . ■

In contrast to Proposition 24, it was proved by Hartung and Nichterlein [12] that deciding whether there is a solution of size k for METRIC DIMENSION is highly unlikely to be solvable by an algorithm of running time $n^{o(k)}$.

Acknowledgments

We thank the anonymous referees for carefully reading the paper, especially for detecting a flaw in the original proof of Theorem 19 and in an erroneous construction that was replaced by Theorems 8 and 13 (which were suggested by one of the referees).

REFERENCES

- [1] R.F. Bailey and P.J. Cameron, Base size, metric dimension and other invariants of groups and graphs, Bull Lond Math Soc 43 (2011), 209–242.
- [2] D.I. Carson, On generalized location domination, Graph Theory, Combinatorics and Applications, Proc 7th Quadrennial International Conference on the Theory and Applications of Graphs, Vol. 1, Kalamazoo, Michigan, USA, 1995, pp. 161–179.
- [3] G. Chartrand, L. Eroh, M. Johnson, and O. Oellermann, Resolvability in graphs and the metric dimension of a graph, Discrete Appl Math 105 (2000), 99–113.

- [4] O. Delmas, S. Gravier, M. Montassier, and A. Parreau, On two variations of identifying codes, *Discrete Math* 311 (2011), 1948–1956.
- [5] F. Foucaud, Combinatorial and algorithmic aspects of identifying codes in graphs, PhD thesis, Université Bordeaux 1, Bordeaux, France, 2012. Available at: <http://tel.archives-ouvertes.fr/tel-00766138>. Accessed on August 21, 2014.
- [6] F. Foucaud, The complexity of the identifying code problem in restricted graph classes, *Proc 24th International Workshop on Combinatorial Algorithms, Lecture Notes in Computer Science*, Vol. 8288, Springer, Rouen, France, 2013, pp. 150–163.
- [7] F. Foucaud, On the decision and approximation complexities for identifying codes and locating-dominating sets in restricted graph classes, Manuscript (2013). Available at: <http://hal.archives-ouvertes.fr/hal-00923356>. Accessed on August 21, 2014.
- [8] S. Gravier, R. Klasing, and J. Moncel, Hardness results and approximation algorithms for identifying codes and locating-dominating codes in graphs, *Algorithmic Oper Res* 3 (2008), 43–50.
- [9] S.L. Hakimi, Optimal distribution of switching centers and the absolute centers and medians of a graph, *Oper Res* 12 (1964), 462–475.
- [10] F. Harary, Status and contrastatus, *Sociometry* 22 (1959), 23–43.
- [11] F. Harary and R.A. Melter, On the metric dimension of a graph, *Ars Combin* 2 (1976), 191–195.
- [12] S. Hartung and A. Nichterlein, On the parameterized and approximation hardness of metric dimension, *Proc 2013 IEEE Conference on Computational Complexity*, Stanford, California, USA, 2013, pp. 266–276.
- [13] T.W. Haynes, M.A. Henning, and J. Howard, Locating and total dominating sets in trees, *Discrete Appl Math* 154 (2006), 1293–1300.
- [14] M.C. Hernando, M. Mora, I.M. Pelayo, C. Seara, and D.R. Wood, Extremal graph theory for metric dimension and diameter, *Electron J Comb* 17 (2010), #R30.
- [15] I. Honkala, T. Laihonen, and S. Ranto, On strongly identifying codes, *Discrete Math* 254 (2002), 191–205.
- [16] D.S. Johnson, Approximation algorithms for combinatorial problems, *J Comput Syst Sci* 9 (1974), 256–278.
- [17] C. Jordan, Sur les assemblages de lignes (in French), *J Reine Angew Math* 70 (1869), 185–190.
- [18] M.G. Karpovsky, K. Chakrabarty, and L.B. Levitin, On a new class of codes for identifying vertices in graphs, *IEEE Trans Inf Theory* 44 (1998), 599–611.
- [19] S. Khuller, B. Raghavachari, and A. Rosenfeld, Landmarks in graphs, *Discrete Appl Math* 70 (1996), 217–229.
- [20] A. Lobstein, Watching systems, identifying, locating-dominating and discriminating codes in graphs: A bibliography, Available at: <http://www.infres.enst.fr/lobstein/debutBIBidetlocdom.pdf>. Accessed on August 21, 2014.
- [21] OEIS Foundation Inc, The On-Line Encyclopedia of Integer Sequences, Available at: <http://oeis.org/A000670>. Accessed on August 21, 2014.
- [22] A. Sebő and E. Tannier, On metric generators of graphs, *Math Oper Res* 29 (2004), 383–393.
- [23] S.J. Seo and P.J. Slater, Open neighborhood locating-dominating sets, *Australas J Combin* 46 (2010), 109–120.
- [24] S.J. Seo and P.J. Slater, Open neighborhood locating-dominating in trees, *Discrete Appl Math* 159 (2011), 484–489.
- [25] P.J. Slater, Leaves of trees, *Congr Numer* 14 (1975), 549–559.
- [26] P.J. Slater, Maximin facility location, *J Res Natl Bur Stand Ser B* 79 (1975), 107–115.
- [27] P.J. Slater, Medians of arbitrary graphs, *J Graph Theory* 4 (1980), 389–392.
- [28] P.J. Slater, Domination and location in acyclic graphs, *Networks* 17 (1987), 55–64.
- [29] P.J. Slater, Dominating and reference sets in a graph, *J Math Phys Sci* 22 (1988), 445–455.
- [30] P.J. Slater, A survey of sequences of central subgraphs, *Networks* 34 (1999), 244–249.
- [31] C. Smart and P.J. Slater, Center, median, and centroid subgraphs, *Networks* 34 (1999), 303–311.
- [32] H.S. Wilf, *Generatingfunctionology*, 3rd edition, A. K. Peters Ltd, Wellesley, MA, 2006.
- [33] B. Zelinka, Medians and peripherians of trees, *Arch Math* 4 (1968), 87–95.