

Locating-Dominating Sets and Identifying Codes in Graphs of Girth at least 5

Camino Balbuena*

Departament de Matemàtica Aplicada III
Universitat Politècnica de Catalunya, Barcelona, Spain
m.camino.balbuena@upc.edu

Florent Foucaud^{†*}

LIMOS - CNRS UMR 6158
Université Blaise Pascal, France
florent.foucaud@gmail.com

Adriana Hansberg^{‡*§}

Instituto de Matemáticas
Universidad Nacional Autónoma de México, México
ahansberg@im.unam.mx

Submitted: Jul 28, 2014; Accepted: Apr 21, 2015; Published: Apr 29, 2015
Mathematics Subject Classification: 05C69

Abstract

Locating-dominating sets and identifying codes are two closely related notions in the area of separating systems. Roughly speaking, they consist in a dominating set of a graph such that every vertex is uniquely identified by its neighbourhood within the dominating set. In this paper, we study the size of a smallest locating-dominating set or identifying code for graphs of girth at least 5 and of given minimum degree. We use the technique of vertex-disjoint paths to provide upper bounds on the minimum size of such sets, and construct graphs who come close to meeting these bounds.

Keywords: Identifying codes; locating-dominating sets; dominating sets; girth; path covers; minimum degree.

*This research was supported by the Ministry of Education and Science, Spain, and the European Regional Development Fund (ERDF) under project MTM2011-28800-C02-02, and under the Catalanian Government project 1298 SGR2009.

[†]Part of this research was done while this author was a postdoctoral researcher at the Departament de Matemàtica Aplicada III of Universitat Politècnica de Catalunya, Barcelona, Spain, at the Department of Mathematics of University of Johannesburg, South Africa, and at LAMSADE - PSL Université Paris-Dauphine, France.

[‡]Research partially supported by the program UNAM-DGAPA-PAPIIT under project IA103915 and by *Fondo Sectorial de Investigación para la Educación* under project 219775 CB-2013-01.

[§]Part of this research was done while this author was a postdoctoral researcher at the Departament de Matemàtica Aplicada III of Universitat Politècnica de Catalunya, Barcelona, Spain.

1 Introduction

Various forms of distinguishing problems in graphs arising from several applications have been studied. Imagine a setting where one wants to detect a hazard in a network (graph) using simple local detectors. Every network node should be within reach of some detector, say at graph distance at most 1: in this case the detectors must form a dominating set. If, in addition, one wants to be able to precisely locate the hazard, every node must be uniquely determined by the set of detectors monitoring (dominating) it. This is the notion of a locating-dominating set or an identifying code (depending on whether the detector nodes should be distinguished themselves).

Since the introduction of locating-dominating sets by Slater [24, 25] and identifying codes by Karpovsky, Chakrabarty and Levitin [13], these concepts have been widely studied and applied to hazard- or fault-detection in networks and facilities [13, 27], routing [16], as well as in relation with graph isomorphism [3] and logical characterizations of graphs [14]. An online bibliography on these topics is maintained by Lobstein [17]. We remark that these problems belong to the more general set of distinguishing or separating problems in graphs and hypergraphs; see the concept of hypergraph *separating systems* [5, 22] (which is also known under the name of *test covers* [7, 19] or *discriminating codes* [8], and is related to a celebrated theorem of Bondy [6]).

In this paper, we study locating-dominating sets and identifying codes in graphs of girth at least 5 (that is, containing no triangle or 4-cycle). Their behaviour in this class is quite different from the class of graphs with girth 3 or 4. We are able to give upper bounds on the smallest size of such sets in terms of the order of the graph, and discuss the tightness of our bounds.

Definitions. All graphs in this paper will be undirected and finite. The order of a graph will be denoted by the letter n . The open and closed neighbourhoods of a vertex x are denoted $N(x)$ and $N[x]$, respectively, and the *degree* of x is the size of its open neighbourhood. A graph is *cubic* if all its vertices have degree 3. A path along vertices x_1, \dots, x_k is denoted $x_1 - \dots - x_k$. The *order* of a path is the number of its vertices, and its *length* is the number of its edges (that is, its order minus one). We may also denote the concatenation of two paths P, P' by $P - P'$. A *Hamiltonian path* of a graph is a path containing all its vertices. Given a set X of vertices in a graph G , $G[X]$ denotes the subgraph of G induced by X . A graph G is *vertex-transitive* if, given any two vertices x and y , there is an automorphism of G mapping x to y .

In a graph G , a vertex *dominates* itself and all its neighbours. A set D of vertices dominates vertex x if some vertex of D dominates x . Similarly, D *2-dominates* x if at least two distinct vertices of D dominate x . Set D is called a *dominating set* if D dominates all vertices in $V(G)$. If a vertex x belongs to the symmetric difference $N[u] \Delta N[v]$ (i.e. x dominates exactly one of u, v) we say that x *separates* u from v .

We have the following definitions of the core concepts of this paper:

Definition 1 ([13, 24, 25]). Given a graph G , a subset C of vertices of $V(G)$ which is both a dominating set and such that all vertex-pairs in $V(G) \setminus C$ are separated by some vertex

of C is called a *locating-dominating set* of G . If *all* vertex-pairs in $V(G)$ are separated by some vertex of C , it is called an *identifying code* of G .

Note that a graph always has a locating-dominating set, but it has an identifying code (it is *identifiable*) if and only if it has no *twins*, i.e. vertices with the same closed neighbourhood. However, for triangle-free graphs (and thus for graphs of girth at least 5), twins cannot have any common neighbour, leading to the following observation:

Observation 2. *A triangle-free graph is identifiable if and only if it has no connected component with two vertices.*

The minimum size of a dominating set, a locating-dominating set, and an identifying code of a graph G are called the *domination number* $\gamma(G)$, the *location-domination number* $\gamma^{\text{LD}}(G)$ and the *identifying code number* $\gamma^{\text{ID}}(G)$ of G , respectively. If G is identifiable we have $\gamma(G) \leq \gamma^{\text{LD}}(G) \leq \gamma^{\text{ID}}(G)$.

Related work. A classic result in domination due to Ore [20] is that for every graph G of order n with minimum degree at least 1, $\gamma(G) \leq \frac{n}{2}$. Later, McCuaig and Shepherd [18] proved that besides seven exceptional graphs, if G is connected and has minimum degree at least 2, then $\gamma(G) \leq \frac{2n}{5}$. For minimum degree at least 3, Reed [21] proved the bound $\gamma(G) \leq \frac{3n}{8}$. More generally, it is known that any graph G with minimum degree δ has domination number $\gamma(G) = O\left(\frac{\log \delta}{\delta}\right)n$ (see [1]), and this bound is asymptotically tight [2]. On the other hand, for connected cubic graphs with $n \geq 9$, Kostochka and Stodolsky [15] proved that $\gamma(G) \leq \frac{4n}{11}$.

A bound of this form does not exist for locating-dominating sets or identifying codes. Indeed, d -regular graphs with locating-dominating number and identifying code number of the form $n\left(1 - \frac{1}{\Theta(d)}\right)$ were constructed by the second author and Perarnau [10]. However, these constructions contain either triangles or 4-cycles, and the same authors showed that for any graph G of order n , girth at least 5 and minimum degree δ , an (asymptotically tight) upper bound of the form $\gamma^{\text{LD}}(G) \leq \gamma^{\text{ID}}(G) = O\left(\frac{\log \delta}{\delta}\right)n$ (similar to the one for dominating sets) holds. However, for small values of δ the bound of [10] is not meaningful; when $\delta = 2$, the second author showed the bound $\gamma^{\text{ID}}(G) \leq \frac{7n}{8} = 0.875n$ in his PhD thesis [9].

In this paper, we study the following question:

Question 3. What are tight upper bounds on $\gamma^{\text{LD}}(G)$ and $\gamma^{\text{ID}}(G)$ for graphs G of given (small) minimum degree $\delta \geq 2$ and girth at least 5?

Our results and structure of the paper. We study the cases where the minimum degree $\delta \in \{2, 3\}$, and also the case of cubic graphs. In Section 2, we give upper bounds on parameters γ^{LD} and γ^{ID} for these graph classes, and discuss their tightness by constructing examples with large values of γ^{LD} and γ^{ID} in Section 3. We briefly conclude in Section 4. A summary of our results is given in Table 1. To obtain the upper bounds, we use the technique of building vertex-disjoint paths of the graph, that was introduced by Reed [21] for dominating sets and was used in related works, see e.g. [15, 28].

	Location-domination number			Identifying code number		
	largest known examples		upper bound	largest known examples		upper bound
	small	arb. large		small	arb. large	
$\delta = 2$	0.5 C_6 [25]	$0.5 - \epsilon$ Prop. 21	0.5 Thm. 8	$\frac{5}{7}$ C_7 [4]	$0.6 - \epsilon$ Prop. 25	$\frac{5}{7} < 0.715$ Thm. 13
$\delta \geq 3$	$\frac{3}{7} > 0.428$ Prop. 24	$\frac{4}{11} - \epsilon > 0.363$ Prop. 23		0.5 Prop. 29	$\frac{5}{11} - \epsilon > 0.454$ Prop. 27	
cubic		$\frac{1}{3}$ Thm. 18 [26]	$\frac{22}{45} < 0.489$ Cor. 12		0.4 Thm. 18 [13]	$\frac{31}{45} < 0.689$ Cor. 17

Table 1: Upper bounds and largest known ratios (in terms of the graph's order) of location-domination and identifying code numbers in connected graphs of girth at least 5 and minimum degree δ .

2 Upper bounds using vertex-disjoint path covers

This section contains the proofs of our upper bounds. We start with some preliminary tools.

2.1 Preliminary lemmas and definitions

Next, we give useful characterizations of locating-dominating sets and identifying codes in graphs of girth 5.

Lemma 4. *Let G be a graph of girth at least 5, and let C be a dominating set of G . Let $X = \{x \in V(G) \setminus C : |N(x) \cap C| = 1\}$. Then C is a locating-dominating set of G if and only if there is an injective function $f : X \rightarrow C$ such that $f(x) \in C \cap N(x)$ for all $x \in X$.*

Proof. If C is a locating-dominating set, $N(x) \cap C \neq N(y) \cap C$ for each pair x, y of vertices of X . Then clearly the function $f : X \rightarrow C$ such that $f(x) = N(x) \cap C$ is injective (if there are two vertices $x, y \in X$ with $y \neq x$ and $f(x) = f(y)$, then x, y would not be separated, a contradiction).

For the sufficiency, suppose that there is an injective function $f : X \rightarrow C$ such that $f(x) \in C \cap N(x)$ for all $x \in X$. Let $Y = V(G) \setminus (C \cup X)$ and let u, v be distinct vertices of $V(G) \setminus C$. If $u, v \in X$, evidently $f(u) \neq f(v)$ and thus $N(u) \cap C \neq N(v) \cap C$. If $u \in X$ and $v \in Y$, then $|N(u) \cap C| = 1$ and $|N(v) \cap C| \geq 2$ and thus $N(u) \cap C \neq N(v) \cap C$. Lastly, if $u, v \in Y$, then $|N(u) \cap C| \geq 2$ and $|N(v) \cap C| \geq 2$. But then $N(u) \cap C \neq N(v) \cap C$ since otherwise there would be a cycle of length 4. Thus, C is a locating-dominating set. \square

Lemma 4 means that in a graph of girth 5, the fact that a dominating set is also locating only depends on the vertices that are dominated by exactly one vertex.

The following is a more complicated version of Lemma 4 for identifying codes. It is a more precise extension of a lemma used by the second author and Perarnau in [10].

Lemma 5. *Let G be an identifiable graph of girth at least 5. Let C be a dominating set of G and let $C_{\geq 3}$ be the set of vertices of C belonging to a connected component of $G[C]$ of size at least 3. Then, C is an identifying code of G if and only if the following conditions hold:*

- (i) *None of the components of $G[C]$ have size 2;*
- (ii) *For $X = \{x \in V(G) \setminus C : |N(x) \cap C| = 1\}$, there is an injective function $f : X \rightarrow C$ such that $f(x) \in C_{\geq 3} \cap N(x)$ for all $x \in X$.*

Proof. First, assume that C is an identifying code of G . Then, Property (i) is clear (otherwise the two vertices of some component C_i of order 2 would not be separated). The proof that Property (ii) holds is similar as for Lemma 4 (by letting $f(x) = N(x) \cap C$ for each $x \in X$). Observe that $f(x) \in C_{\geq 3}$, otherwise x and $f(x)$ would not be separated.

For the other side, assume that C is a dominating set fulfilling Properties (i) and (ii) and, by contradiction, assume that there are two distinct vertices x, y that are not separated by C , i.e. $N[x] \cap C = N[y] \cap C$.

Assume first that x and y are adjacent. As $N[x] \cap C = N[y] \cap C \neq \emptyset$, it follows that $N[x] \cap C = N[y] \cap C \subseteq \{x, y\}$ (since there is no triangle in G). If both x, y belong to C , x and y induce a component of $G[C]$ of size 2, a contradiction. Otherwise, exactly one of them belongs to C (say x). But then y is only dominated by x , which does not belong to $C_{\geq 3}$, a contradiction to Property (ii).

Thus, x and y are non-adjacent and, since there are no 4-cycles, $|N(x) \cap N(y)| \leq 1$. Hence, there is a vertex z with $N(x) \cap C = N(y) \cap C = \{z\}$. It follows that $x, y \in X$ but $f(x) = z = f(y)$, a contradiction. \square

We now define the key concept of vertex-disjoint path cover of a graph, and introduce some related notation.

Definition 6. A *vertex-disjoint path cover* (*vdp-cover* for short) of G is a partition of $V(G)$ into sets of vertices, each of them inducing a graph with a Hamiltonian path.

For $0 \leq i \leq 4$, a path whose order is congruent to i modulo 5 is called an $(i \bmod 5)$ -*path*, and a path of order j is a j -*path* (an empty path is a 0-path). Given a vdp-cover \mathcal{S} , we will denote by \mathcal{S}_i the set of $(i \bmod 5)$ -paths in \mathcal{S} , and by \mathcal{T}_i , the set of i -paths in \mathcal{S} .

The following result of Reed [21] will be used.

Theorem 7 ([21]). *Every connected cubic graph of order n has a vdp-cover with at most $\frac{n}{9}$ sets.*

2.2 Locating-dominating sets

The bound given in the following theorem also follows from a stronger result in a recent paper by Garijo, González and Márquez [11] (see there Proposition 6.6). However, we give an independent proof by a completely different method, which is a good and simple illustration of this technique that will be used several times in this paper.

Theorem 8. *Let G be a graph of order n , girth at least 5 and minimum degree at least 2. Then $\gamma^{LD}(G) \leq \frac{n}{2}$.*

Proof. Let \mathcal{S} be a vdp-cover of G and let \mathcal{T}_1 and \mathcal{T}_3 be the sets of paths of order 1 and 3 in \mathcal{S} , respectively. Let \mathcal{S} be chosen such that $2|\mathcal{T}_1| + |\mathcal{T}_3|$ is minimized. Without loss of generality, we can assume that all paths in \mathcal{S} have length at most 5, since otherwise we can split any longer path into paths of lengths 2 or 5 without affecting the minimality condition. For each path $P \in \mathcal{S}$ of length $1 \leq r \leq 5$, we define an order $P = x_0 - x_1 - \dots - x_{r-1}$ with x_i adjacent to x_{i+1} for $0 \leq i < r - 1$. Let D be the set of vertices containing all vertices of the paths of \mathcal{S} of odd index (i.e. all x_1 's and all x_3 's). Note that D clearly dominates all vertices, except possibly the vertices in a 1-path. Also, we define a function f on all vertices with index 0 or 4 in the following way. If $P = x_0 - x_1 - \dots - x_{r-1}$ is an r -path with $2 \leq r \leq 5$, then $f(x_0) = x_1$ and, if $r = 5$, $f(x_4) = x_3$. According to Lemma 4, if the end-vertices of the 3-paths in \mathcal{S} have, besides of their neighbour on the path, a second neighbour in D and if all vertices of the 1-paths from \mathcal{S} have two neighbours in D , then the restriction of f to the set of 1-dominated vertices is injective and therefore D is a locating-dominating set.

We will show that every vertex of a 1-path and every end-vertex of a 3-path has no neighbour outside of D ; which by the previous discussion suffices for D being a locating-dominating set. Herefor, we say that a vertex x is a (p, q) -vertex if it belongs to a path P of order $p + q + 1$ of \mathcal{S} and the two paths obtained from P by removing x have orders p and q . Observe that a (p, q) -vertex is the same as a (q, p) -vertex. Further, we say that, for fixed p and q , the (p, q) -vertices are *good* if they all belong to D , otherwise they are *bad*. Taking into account that $p + q + 1 \leq 5$, we have the following pairs (p, q) such that (p, q) -vertices are bad: $(0, 0)$, $(0, 1)$, $(0, 2)$, $(0, 3)$, $(0, 4)$, $(1, 2)$ and $(2, 2)$.

Let $P = x \in \mathcal{T}_1$ be a 1-path. If x is adjacent to a $(0, q)$ -vertex for some $q \in \{0, 1, 2, 3, 4\}$, then we can replace the 1-path and the $(q + 1)$ -path by a $(q + 2)$ -path, obtaining in all cases a lower value for the sum $2|\mathcal{T}_1| + |\mathcal{T}_3|$, a contradiction. Hence suppose that x is adjacent to either a $(1, 2)$ -vertex or to a $(2, 2)$ -vertex. Then we can substitute the 1-path and the 4- or 5-path by a 2-path and a 3- or 4-path, obtaining in each case a lower value for the sum $2|\mathcal{T}_1| + |\mathcal{T}_3|$, which is a contradiction. Hence, x has to be adjacent only to good vertices. As $\delta(G) \geq 2$, it follows that x is adjacent to two vertices from D . Completely analogous we obtain a contradiction when P is a 3-path having an end-vertex adjacent to a bad vertex. Altogether, it follows that all vertices not in D have either an assignment via f or two neighbours in D . Hence, by Lemma 4, D is a locating-dominating set. Since each path from \mathcal{S} has at most half of its vertices in D , we obtain $\gamma^{\text{LD}}(G) \leq |D| \leq \frac{n}{2}$. \square

Theorem 8 is tight for the cycles C_6 and C_8 , which can easily be seen to have location-domination numbers 3 and 4, respectively (see also [25]). In Proposition 21, we will give a construction of arbitrarily large connected graphs based on copies of C_6 .

Next, given a vdp-cover \mathcal{S} of a graph G with girth 5, we will show how to construct a set $D(\mathcal{S})$ and an injective function $f : X \rightarrow D(\mathcal{S})$ (where X is the set of 1-dominated vertices of $V(G) \setminus D(\mathcal{S})$) meeting the conditions of Lemma 4. We will build $D(\mathcal{S})$ by taking roughly two vertices out of five in each path of \mathcal{S} , then adding a few vertices for each path whose length is nonzero modulo 5.

Definition 9. Let G be a graph of girth at least 5 and \mathcal{S} be a vdp-cover of G . Then, the set $D(\mathcal{S})$ and the function $f_{D(\mathcal{S})}$ are constructed as follows.

For each path $P = x_0 - \dots - x_{p-1}$ in \mathcal{S} , we do the following. Assume that $P \in \mathcal{S}_i$ ($0 \leq i \leq 4$), that is, $p = 5k + i$ for some $k \geq 0$. If $k \geq 1$, $D(\mathcal{S})$ contains the set $\{x_j \in V(P), j = 1, 3 \pmod 5, j < 5k\}$.

Now, if $k \geq 0$ and P belongs to $\mathcal{S} \setminus \mathcal{S}_0$, we add some vertices to $D(\mathcal{S})$ according to the following case distinction:

- If $P \in \mathcal{S}_1$, we let $D(\mathcal{S})$ contain x_{p-1} .
- If $P \in \mathcal{S}_2$, $D(\mathcal{S})$ also contains x_{p-2} and $f_{D(\mathcal{S})}(x_{p-1}) = x_{p-2}$.
- If $P \in \mathcal{S}_3$, $D(\mathcal{S})$ also contains $\{x_{p-3}, x_{p-2}\}$ and $f_{D(\mathcal{S})}(x_{p-1}) = x_{p-2}$.
- If $P \in \mathcal{S}_4$, $D(\mathcal{S})$ also contains $\{x_{p-3}, x_{p-1}\}$ and $f_{D(\mathcal{S})}(x_{p-4}) = x_{p-3}$.

To finish the construction of the function $f_{D(\mathcal{S})}$, for $j < 5k$, if $x_j \notin D(\mathcal{S})$ and $j = 0 \pmod 5$, $f_{D(\mathcal{S})}(x_j) = x_{j+1}$; if $j = 4 \pmod 5$, $f_{D(\mathcal{S})}(x_j) = x_{j-1}$.

An illustration of Definition 9 is given in Figure 1.




Figure 1: Illustration of set $D(\mathcal{S})$.

Lemma 10. Let G be a graph of girth at least 5 having a vdp-cover \mathcal{S} . Then $D(\mathcal{S})$ is a locating-dominating set of G .

Proof. The proof follows from Lemma 4; indeed, each vertex x of a path $P \in \mathcal{S}$ and $x \notin D(\mathcal{S})$ that is not 2-dominated has an image $f_{D(\mathcal{S})}(x) \in P$ (and no other such vertex y has $f_{D(\mathcal{S})}(x) = f_{D(\mathcal{S})}(y)$). It follows that the restriction of $f_{D(\mathcal{S})}$ to the set X of 1-dominated vertices is injective. \square

Now, using Theorem 7 and the above construction of the set $D(\mathcal{S})$, we can give an improved bound for cubic graphs, based on the following general theorem:

Theorem 11. Let G be a graph of order n , girth at least 5 and having a vdp-cover with $\alpha \cdot n$ paths. Then $\gamma^{LD}(G) \leq \frac{2+4\alpha}{5}n$.

Proof. Let \mathcal{S} be a vdp-cover of G of size at most $\alpha \cdot n$. We consider the set $D(\mathcal{S})$ defined in Definition 9. By Lemma 10, $D(\mathcal{S})$ is a locating-dominating set of G . It remains to estimate the size of $D(\mathcal{S})$.

For each path P in \mathcal{S}_i with $5k + i$ vertices ($k \geq 0$) blue, we have added $\frac{2k}{5}$ vertices of P to $D(\mathcal{S})$ in the first step of the construction. Then, in the second step, for each path in $\mathcal{S}_1 \cup \mathcal{S}_2$ and $\mathcal{S}_3 \cup \mathcal{S}_4$, we have added one and two additional vertices, respectively. So in total we get:

$$\begin{aligned} |D(\mathcal{S})| &\leq \frac{2}{5}(n - |\mathcal{S}_1| - 2|\mathcal{S}_2| - 3|\mathcal{S}_3| - 4|\mathcal{S}_4|) + |\mathcal{S}_1| + |\mathcal{S}_2| + 2|\mathcal{S}_3| + 2|\mathcal{S}_4| \\ &= \frac{2}{5}n + \frac{3}{5}|\mathcal{S}_1| + \frac{1}{5}|\mathcal{S}_2| + \frac{4}{5}|\mathcal{S}_3| + \frac{2}{5}|\mathcal{S}_4| \\ &\leq \frac{2}{5}n + \frac{4}{5}|\mathcal{S}| \\ &\leq \frac{2 + 4\alpha}{5}n. \end{aligned}$$

□

We get the following corollary of Theorems 7 and 11:

Corollary 12. *Let G be a connected cubic graph of order n and girth at least 5. Then $\gamma^{LD}(G) \leq \frac{22}{45}n < 0.489n$.*

2.3 Identifying codes

The methods used in this subsection are similar to the ones of Subsection 2.2, but the proofs are more intricate.

Theorem 13. *Let G be an identifiable graph of order n , girth at least 5 and minimum degree $\delta \geq 2$. Then, $\gamma^{ID}(G) \leq \frac{5}{7}n < 0.715n$.*

Proof. Given a vdp-cover \mathcal{S} of G , let \mathcal{T}_i be the set of paths of order exactly i of \mathcal{S} (in this proof we do not consider the orders modulo 5). We choose \mathcal{S} such that

$$4|\mathcal{T}_1 \cup \mathcal{T}_4| + 3|\mathcal{T}_2 \cup \mathcal{T}_3| + 2|\mathcal{T}_8 \cup \mathcal{T}_9| \tag{1}$$

is minimized. Let $P \in \mathcal{S}$ be an r -path with $r \geq 10$. Then we can replace P by paths of orders 5, 6 and 7 without affecting the minimality of (1):

- If $r \equiv 0 \pmod{5}$, then we can replace P by 5-paths.
- If $r \equiv 1 \pmod{5}$, then we can replace P by one 6-path and the remaining part by 5-paths.
- If $r \equiv 2 \pmod{5}$, then we can replace P by one 7-path and the remaining part by 5-paths.

- If $r \equiv 3 \pmod{5}$, then we can replace P by one 6-path, one 7-path, and the remaining part by 5-paths.
- If $r \equiv 4 \pmod{5}$, then we can replace P by two 7-paths and the remaining part by 5-paths.

Hence, without loss of generality, we can assume that there are no paths of length 10 or more in \mathcal{S} . Now, we define a set C in the following way. For each r -path $P = x_0 - x_1 - \dots - x_{r-1}$ of \mathcal{S} , we add some vertices to C and define a function f according to the following distinction:

- If $r = 2$, then let C contain x_1 .
- If $r = 3$, then let C contain x_1 and x_2 .
- If $r = 4$, then let C contain x_0 and x_3 ; let $f(x_1) = x_0$ and $f(x_2) = x_3$.
- If $5 \leq r \leq 7$, then let C contain x_1, x_2, \dots, x_{r-2} ; let $f(x_0) = x_1$ and $f(x_{r-1}) = x_{r-2}$.
- If $r = 8$, then let C contain x_1, x_2, x_3, x_6 and x_7 ; let $f(x_4) = x_3$ and $f(x_5) = x_6$.
- If $r = 9$, then let C contain x_1, x_2, x_3, x_4, x_7 and x_8 ; let $f(x_5) = x_4$ and $f(x_6) = x_7$.

An illustration of set C is given in Figure 2. We will show that C is an identifying code of G .




Figure 2: Illustration of set C in the proof of Theorem 13.

As in the proof of Theorem 8, we say that a vertex x is a (p, q) -vertex if it belongs to a path P of order $p + q + 1$ of \mathcal{S} and the two paths obtained from P by removing x have orders p and q . Observe that a (p, q) -vertex is the same as a (q, p) -vertex. Further, we

say that, for fixed p and q , the (p, q) -vertices are *good* if they all belong to C , otherwise they are *bad*. Taking into account that $p + q + 1 \leq 9$, we have the following set B of pairs (p, q) such that (p, q) -vertices are bad:

$$B = \{(0, 0), (0, 1), (0, 2), (0, 4), (0, 5), (0, 6), (0, 7), (0, 8), (1, 2), (3, 4), (2, 5), (2, 6), (3, 5)\}.$$

Now we will prove the following claims.

Claim 13.A. *For a path $P \in \mathcal{S}$ of order $r \in \{8, 9\}$, we can assume that the end-vertex x_{r-1} , which belongs to C , has either a second neighbour in P contained in C (i.e. different from its predecessor x_{r-2} in P) or it has a neighbour outside P .*

Let $r = 8$ and $P = x_0 - x_1 - \dots - x_7$ and, following the construction of C , we have $x_1, x_2, x_3, x_6, x_7 \in C$. By contradiction, suppose that x_7 is not adjacent to any of x_1, x_2, x_3 . Suppose also that x_7 has no neighbour outside P . Since G has girth at least 5, x_7 is neither adjacent to x_4 nor to x_5 . Hence, as $\delta \geq 2$, x_7 has to be adjacent to x_0 . Now, either $G = C_8$ or one of the vertices from P has one neighbour outside P . In the first case, an independent set of size 4 is an identifying code of $G = C_8$ and satisfies the desired bound. Hence we may assume that $G \neq C_8$ and thus there is a vertex from P having a neighbour outside P . In this case, we may reorder the vertices along the cycle such that x_7 has one neighbour outside P . Hence, Claim 13.A follows for $r = 8$. The same argument can be used to prove the case $r = 9$.

Claim 13.B. *Let $r \in \{1, 2, 3, 4, 8, 9\}$ and let x be an end-vertex of an r -path $P \in \mathcal{S}$. Then all neighbours of x outside P are good vertices.*

Suppose that, for some $r \in \{1, 2, 3, 4, 8, 9\}$, there is an end-vertex of an r -path P which is adjacent to a (p, q) -vertex in $P' \in \mathcal{S}$ with $P \neq P'$ and $(p, q) \in B$. Note that we can replace P and P' by either an $(r + p + 1)$ -path and a q -path or by a p -path and an $(r + q + 1)$ -path.¹ We will see that, in each case, we obtain a vdp-cover which contradicts the minimality of (1). If $p = 0$, then we can join the r -path together with the $(q + 1)$ -path obtaining an $(r + q + 1)$ -path. This gives in all cases a lower value for the sum (1), which is a contradiction. Hence we can suppose that $(p, q) \in \{(1, 2), (3, 4), (2, 5), (2, 6), (3, 5)\}$. When $r = 4$ and (p, q) is arbitrary or when $r = 2$ and $(p, q) = (1, 2)$, we can replace the r - and the $(p + q + 1)$ -path by an $(r + q + 1)$ - and a p -path and we obtain in all cases a lower value for (1). For $r \in \{1, 3, 8, 9\}$ and (p, q) is arbitrary or $r = 2$ and $(p, q) \neq (1, 2)$, we can replace the r - and the $(p + q + 1)$ -paths by an $(r + p + 1)$ - and a q -path and we obtain always a lower value for (1). Since we obtain in all cases a contradiction to the minimality of (1), it follows that, for $r = 1, 2, 3, 4, 8, 9$, every end-vertex of an r -path is adjacent to a good vertex, proving Claim 13.B.

Claim 13.C. *Every vertex from a 1-path is adjacent to two vertices of C .*

As $\delta \geq 2$ and since by Claim 13.B, the vertex of a 1-path cannot be adjacent to a bad vertex, then it has to be adjacent to at least two good vertices, proving Claim 13.C.

¹Whenever we consider a new s -path with $s \geq 10$, we implicitly assume that, as done in the beginning of the proof, it is cut into smaller paths.

Claim 13.D. *There are no components of $G[C]$ that have size at most 2.*

Since the girth of G is at least 5 and $\delta \geq 2$, all end-vertices of a 2-, 3- or 4-path P have a neighbour outside P . By Claim 13.B, these neighbours have to be good vertices. Hence, there are no 1-components in $G[C]$. On the other side, if P is an 8- or a 9-path, Claim 13.A implies that the end-vertices of P have either a further neighbour in P belonging to C or they have a neighbour outside P , which, by Claim 13.B, is a good vertex. Thus, the only possibilities to have 2-components in $G[C]$ are given when two 2-paths or one 2-path and one 4-path or two 4-paths are connected through their good end-vertices. In these cases we could transform them into a 4-path, a 6-path or an 8-path which would contribute less to the sum (1) than the original paths, which is a contradiction, proving Claim 13.D.

Hence, by Claims 13.A, 13.B, 13.C and 13.D and by the construction of the function f , C fulfils the conditions of Lemma 5, which certifies that it is an identifying code. Since at most $\frac{5}{7}|P|$ vertices from every path $P \in \mathcal{S}$ belong to C , C is an identifying code of G of cardinality at most $\frac{5}{7}n$. \square

Theorem 13 is tight for the cycle C_7 , which can easily be seen to have identifying code number 5 (see also [4]).

As for locating-dominating sets, given a vdp-cover \mathcal{S} of a graph G with girth 5, we define a set $C(\mathcal{S})$ and a function $f_{C(\mathcal{S})}$ as follows.

Definition 14. Let G be a graph of girth at least 5 and \mathcal{S} be a vdp-cover of G . Then, the set $C(\mathcal{S})$ and the function $f = f_{C(\mathcal{S})}$ are constructed as follows.

For each path $P = x_0 - \dots - x_{p-1}$ of \mathcal{S} , we do the following. Assume that $P \in \mathcal{S}_i$ ($0 \leq i \leq 4$), that is, $p = 5k + i$ for some $k \geq 0$. If $k \geq 1$, $C(\mathcal{S})$ contains the set $\{x_j \in V(P), j = 1, 2, 3 \bmod 5, j < 5k\}$.

Now, for $k \geq 0$, if P belongs to $\mathcal{S} \setminus \mathcal{S}_0$, we add some vertices to $C(\mathcal{S})$ according to the following case distinction:

- If $P \in \mathcal{S}_1$ and $k \geq 1$, we let $C(\mathcal{S})$ contain x_{p-2} and $f(x_{p-1}) = x_{p-2}$. If $k = 0$, $C(\mathcal{S})$ contains x_0 .
- If $P \in \mathcal{S}_2$ and $k \geq 1$, $C(\mathcal{S})$ also contains $\{x_{p-3}, x_{p-2}\}$ and $f(x_{p-1}) = x_{p-2}$. If $k = 0$, $C(\mathcal{S})$ contains $\{x_0, x_1\}$.
- If $P \in \mathcal{S}_3$ and $k \geq 1$, $C(\mathcal{S})$ also contains $\{x_{p-3}, x_{p-2}, x_{p-1}\}$. If $k = 0$, $C(\mathcal{S})$ contains $\{x_0, x_1, x_2\}$.
- If $P \in \mathcal{S}_4$ and $k \geq 1$, $C(\mathcal{S})$ also contains $\{x_{p-4}, x_{p-3}, x_{p-2}\}$ and $f(x_{p-1}) = x_{p-2}$. If $k = 0$, $C(\mathcal{S})$ contains $\{x_0, x_1, x_2\}$ and $f(x_3) = x_2$.

To finish the construction of the function f , for $j < 5k$, if $x_j \notin C(\mathcal{S})$ and $j = 0 \bmod 5$, $f(x_j) = x_{j+1}$; if $j = 4 \bmod 5$, $f(x_j) = x_{j-1}$. Note that each vertex $x \in P$ of $V(G) \setminus C(\mathcal{S})$ has an image $f(x)$ belonging to P .

An illustration of Definition 14 is given in Figure 3.




Figure 3: Illustration of set $C(\mathcal{S})$.

Lemma 15. *Let G be an identifiable graph of girth at least 5 having a vdp-cover \mathcal{S} . Then $C(\mathcal{S})$ is a dominating set, and all pairs of vertices are separated, except possibly pairs x, y of vertices such that $x - y$ forms a path of \mathcal{S} .*

Proof. The proof follows from Lemma 5; indeed, each vertex x of $V(G) \setminus C(\mathcal{S})$, with $x \in P$ and $P \in \mathcal{S}$, that is not 2-dominated has an image $f_{C(\mathcal{S})}(x) \in P$, the restriction of $f_{C(\mathcal{S})}$ to the set of 1-dominated vertices is injective, and the only potentially isolated vertices in $C(\mathcal{S})$ are vertices v belonging to a path of \mathcal{S} of order 1 (hence by construction no vertex x has $f_{C(\mathcal{S})}(x) = v$). \square

Similarly to Theorem 11 for locating-dominating sets, we have the following generic theorem:

Theorem 16. *Let G be an identifiable graph of order n , girth at least 5 and having a vdp-cover with $\alpha \cdot n$ paths. Then $\gamma^{LD}(G) \leq \frac{3+4\alpha}{5}n$.*

Proof. Let \mathcal{S} be the vdp-cover of G . The idea is to construct a set C and an injective function $f : X \rightarrow C$ (where X is the set of 1-dominated vertices of $V(G) \setminus C$) meeting the conditions of Lemma 5. We will build C by taking roughly three vertices out of five in each path of \mathcal{S} , then adding a few vertices for each path whose length is nonzero modulo five, and finally performing a few local modifications.

Step 1: Constructing an initial pseudo-code. We construct $C = C(\mathcal{S})$ and $f = f_{C(\mathcal{S})}$ by the procedure described in Definition 14.

Step 2: Taking care of components of $G[C]$ of order 2. By Lemma 15, all conditions of Lemma 5 (where we consider the restriction of f to the set X of 1-dominated vertices) are fulfilled, except for Property (i): there might be some paths in \mathcal{S}_2 of order exactly 2 and forming a connected component of $G[C]$ (second item of our case distinction). Let P be such a path, and $V(P) = \{x_0, x_1\}$. Then, since G is identifiable, one of x_0, x_1 (say x_1) has a neighbour y , and since P is a connected component in $G[C]$, $y \notin C$. By the construction of C , y belongs to a path and is adjacent to vertex $f(y)$ in C . We perform the following modification: remove x_0 from C , put y instead, and let $f(x_0) = x_1$. It is

clear that repeating this for each such case, we get rid of all components of order 2 in $G[C]$.

Now, all conditions of Lemma 5 are fulfilled, hence C is an identifying code of G .

Step 3: Saving one vertex for each path of \mathcal{S}_3 .

We consider all paths in \mathcal{S}_3 one by one, in an arbitrary order. For each such path P with $V(P) = \{x_0, \dots, x_{p-1}\}$ ($p = 5k + 3$ for some $k \geq 0$), we remove $x_{p-3}, x_{p-2}, x_{p-1}$ from C . We now distinguish some cases.

If $x_i \in \{x_{p-2}, x_{p-1}\}$ has a neighbour in C , then, we add both x_{p-2}, x_{p-1} to C and let $f(x_{p-3}) = x_{p-2}$. Similarly, if x_{p-3} has a neighbour in C , we add both x_{p-3}, x_{p-2} to C and let $f(x_{p-1}) = x_{p-2}$. Note that in both cases, the two new code-vertices are now part of a component of $G[C]$ of order at least 3, hence all conditions of Lemma 5 are preserved.

If none of $x_{p-3}, x_{p-2}, x_{p-1}$ have a neighbour in C , we add x_{p-3} and x_{p-1} to C . Note that x_{p-2} is now 2-dominated, hence all conditions of Lemma 5 are again preserved.

Repeating this at every step, C is still an identifying code, and we have decreased the size of C by $|\mathcal{S}_3|$.

Step 4: Estimating the size of the code. It remains to compute the size of C .

For each path P in \mathcal{S}_i with $5k + i$ vertices, we have added $\frac{3k}{5}$ vertices of P to C in the first phase of the construction of $C(\mathcal{S})$ (Definition 14). Then, in the second phase of the construction of Definition 14, for each path in $\mathcal{S}_1, \mathcal{S}_2, \mathcal{S}_3, \mathcal{S}_4$, we have added one, two, three and three additional vertices, respectively. In Steps 2 and 3, we did not change the size of C , but in Step 4, we were able to reduce the size of C by one for each path in \mathcal{S}_3 . So in total we have:

$$\begin{aligned} |C| &\leq \frac{3}{5}(n - |\mathcal{S}_1| - 2|\mathcal{S}_2| - 3|\mathcal{S}_3| - 4|\mathcal{S}_4|) + |\mathcal{S}_1| + 2|\mathcal{S}_2| + 3|\mathcal{S}_3| + 3|\mathcal{S}_4| - |\mathcal{S}_3| \\ &= \frac{3}{5}n + \frac{2}{5}|\mathcal{S}_1| + \frac{4}{5}|\mathcal{S}_2| + \frac{1}{5}|\mathcal{S}_3| + \frac{3}{5}|\mathcal{S}_4| \\ &\leq \frac{3}{5}n + \frac{4}{5}|\mathcal{S}| \\ &\leq \frac{3 + 4\alpha}{5}n. \end{aligned}$$

□

We get the following improvement for cubic graphs, a corollary of Theorems 7 and 16:

Corollary 17. *Let G be a connected cubic identifiable graph of order n and girth at least 5. Then $\gamma^{\text{ID}}(G) \leq \frac{31}{45}n < 0.689n$.*

3 Constructions

In this section, we provide constructions of connected graphs with girth at least 5 and large location-domination or identifying code number. First of all, the following result is a lower bound on parameters γ^{LD} and γ^{ID} depending on the maximum degree Δ of a graph. It will be useful since it also applies to Δ -regular graphs.

Theorem 18 ([9, 13, 26]). *Let G be a graph of order n and maximum degree Δ . Then $\gamma^{LD}(G) \geq \frac{2n}{\Delta+3}$. If G is identifiable, then $\gamma^{LD}(G) \geq \frac{2n}{\Delta+2}$, and any identifying code of this size is an independent 2-dominating set whose vertices all have degree Δ in G .*

We remark that the last part of the statement is not very difficult to obtain from the proof of the bound; a proof is available in the first author's PhD thesis [9, Section 4.1].

3.1 Generic constructions

We now define constructions based on the Petersen graph that will be used later on.

Definition 19. Denote by P_{10} the Petersen graph with $V(P_{10}) = \{0, 1, 2, 3, 4, 5, 6, 7, 8, 9\}$, and $0-1-2-\dots-9$ one of its Hamiltonian paths, such that vertices 1 and 9 are adjacent. Let P_{11} be the graph obtained from P_{10} by subdividing once the edge $\{0, 1\}$, and calling the new vertex x .

The graphs of Definition 19 are illustrated in Figure 4.




Figure 4: The Petersen graph P_{10} and its modification P_{11} . The black vertices form an optimal identifying code and locating-dominating set of P_{10} .

Definition 20. For any $k \geq 2$, let G_{11}^k be the graph formed by a vertex y connected to k copies of P_{11} (each attached via vertex x).

The graph G_{11}^k is illustrated in Figure 5.

3.2 Locating-dominating sets

We now give constructions with large location-domination number. The first construction is based on copies of the 6-cycle C_6 .

Proposition 21. *There are infinitely many connected graphs G of order n , girth 5 and minimum degree 2 with $\gamma^{LD}(G) = \frac{n-1}{2}$.*




Figure 5: The graph G_{11}^k .

Proof. Consider the graph G obtained from one vertex x and $k \geq 2$ disjoint copies of C_6 , each joined to x by exactly one edge. We have $n = 6k + 1$, and we claim that $\gamma^{\text{LD}}(G) = 3k$. It is easy to check that a set consisting of three vertices in each copy of C_6 (see Figure 6) is locating-dominating. For the lower bound, assume that D is an optimal locating-dominating set, and that $x \notin D$. Then, each copy of C_6 contains at least $\gamma^{\text{LD}}(C_6) = 3$ vertices of D , and we are done. Hence, assume that $x \in D$. Each copy of C_6 has at least two vertices from D (otherwise D is not dominating). Assume some copy contains exactly two (y, z) : then the neighbour of x in that copy must be only dominated by x . Indeed, if this is not the case (say he is dominated also by y), there would be two vertices in this copy of C_6 that are not in D but only dominated by z , a contradiction. But now observe that in the whole graph, at most one vertex of $V(G) \setminus D$ can be dominated only by x , hence all other copies of C_6 contain three vertices of D , and we are done. \square




Figure 6: A family of connected graphs with location-domination number $\frac{n-1}{2}$.

We will use the following lemma about the graph P_{11} .

Lemma 22. *Let G be a graph of girth 5 containing a copy P of P_{11} as an induced subgraph, such that in P , only vertex x has neighbours out of P . Let D be a locating-dominating set of G . Then, we have $|D \cap V(P)| \geq 4$.*

Proof. By contradiction, we assume that $D_P = D \cap V(P)$ has size 3. If $x \notin D_P$, then D_P must form a locating-dominating set of $P \setminus \{x\}$. By Theorem 18, $\gamma^{LD}(G) \geq \frac{20}{6} > 3$, a contradiction. Hence, $x \in D_P$. But now it is not possible to even dominate the remaining vertices with just two vertices, a contradiction. \square

Proposition 23. *There are infinitely many connected graphs G of order n , girth 5 and minimum degree 3 with $\gamma^{LD}(G) = \frac{4}{11}(n - 1) > 0.363n$.*

Proof. Consider the graph G_{11}^k ($k \geq 3$) from Definition 20, which has $n = 11k + 1$ vertices. A locating-dominating set of size $4k$ is given by selecting vertices $\{x, 3, 6, 9\}$ of each copy of P_{11} (see Figure 5(a)). By Lemma 22, this is optimal. \square

The Heawood graph H_{14} is a well-known Hamiltonian cubic vertex-transitive graph on 14 vertices and with girth 6. Given its vertex set $\{0, 1, \dots, 13\}$, its edges are given by a Hamiltonian cycle $0 - 1 - 2 - \dots - 13$ and $\{0, 5\}$, $\{1, 10\}$, $\{2, 7\}$, $\{3, 12\}$, $\{4, 9\}$, $\{6, 11\}$ and $\{8, 13\}$. See Figure 7 for an illustration.




Figure 7: The Heawood graph with a minimum locating-dominating set (black vertices).

Proposition 24. *The Heawood graph H_{14} has $\gamma^{LD}(H_{14}) = 6 = \frac{3}{7}n > 0.428n$.*

Proof. A locating-dominating set of size 6 is for example $\{1, 4, 6, 8, 10, 13\}$.

We now prove that no locating-dominating set of size 5 exists. Assume by contradiction that there is a locating-dominating set D of H_{14} of size 5. Let $m(D)$ and $m(D, S)$ count the number of edges between vertices of D and the edges between D and $S = V(H_{14}) \setminus D$, respectively. Since at most $|D|$ vertices from S can be dominated by a single vertex of D , we have $m(D, S) \geq |D| + 2(|S| - |D|) = 13$. On the other hand, since H_{14} is cubic, $m(D, S) = 15 - 2m(D)$. Hence, we have $m(D) \leq 1$.

Therefore, we have at least three vertices in D that are adjacent only to vertices of S . Since H_{14} is vertex-transitive, we assume without loss of generality that vertex 0 is such a vertex. Among the neighbours of 0 (vertices 1, 5, 13), at most one is dominated only by 0.

Assume that one of them is in that case. By the symmetries of the graph, there are automorphisms pairwise exchanging edges $\{0, 1\}$, $\{0, 5\}$, $\{0, 13\}$. Hence, without loss of

generality, we can assume that vertex 5 is 1-dominated, but vertices 1, 13 are 2-dominated. Hence, vertices 4, 6 $\notin D$ but at least one vertex among 2, 10 and 8, 12 belongs to D , respectively. Moreover, in order to dominate vertices 4 and 6, one of 3, 9 and 7, 11 belongs to D , respectively. Since these four sets are disjoint and $|D| = 5$, D contains *exactly* one of each.

We first assume that $2 \in D$: hence $10 \notin D$. If also $7 \in D$ (and $11 \notin D$), both 9, 12 $\in D$ in order to dominate 10 and 11, respectively. Then $D = \{0, 2, 7, 9, 12\}$ but 4, 10 are both dominated only by 9, a contradiction. Hence, $7 \notin D$ and $11 \in D$. Then, 9 $\in D$ in order to separate 6, 10; then, 3 $\notin D$ and 12 $\in D$, otherwise 6, 12 are not separated. Hence $D = \{0, 2, 9, 11, 12\}$ but 4, 8 are both dominated only by 9, a contradiction.

Hence, $2 \notin D$ and $10 \in D$. If 3 $\notin D$, then 9 $\in D$ and moreover 12 $\in D$ in order to dominate 3 (hence 8 $\notin D$). Since 7 is dominated, 7 is the last vertex of D . But then 2, 6 are both dominated only by 7, a contradiction. Hence, 3 $\in D$ and 9 $\notin D$. To separate 2, 4, 7 $\in D$ (hence 11 $\notin D$). Then 8 is the last vertex of D , otherwise it would not be separated by 6. But then 4, 12 are not separated, a contradiction.

Therefore, we can assume that all neighbours of 0 are 2-dominated. Hence, at least one vertex among $\{2, 10\}$, $\{4, 6\}$ and $\{8, 12\}$, respectively, belongs to D . Assume first that $2 \in D$. Then, in order for 10 to be dominated, one of 9, 10, 11 belongs to D . Then, exactly one of 8, 12 belongs to D . If $10 \in D$, one of 8, 12 would not be dominated, a contradiction. If 9 $\in D$, then 12 $\in D$ (otherwise it is not dominated). But then, both 8, 10 are only dominated by 9, a contradiction. A similar contradiction follows if $11 \in D$.

Hence, $2 \notin D$, and $10 \in D$. Then, (exactly) one of 3, 7 belongs to D , otherwise 2 is not dominated. If 3 $\in D$ and 7 $\notin D$, 8 $\in D$ (otherwise 8 is not dominated). Since 6 must be dominated, 6 itself is the last vertex of D ; but then, 4, 12 are both only dominated by 3, a contradiction. Hence, If 7 $\in D$ and 3 $\notin D$. Then, 12 $\in D$ (otherwise it is not dominated). Hence, 8 $\notin D$. But now, both 2, 8 are only dominated by 7, a contradiction.

Therefore, D does not exist, which completes the proof. \square

3.3 Identifying codes

We now give constructions with large identifying code number. We start with a construction based on the 5-cycle C_5 , which has identifying code number 3 [4].

Proposition 25. *There are infinitely many connected graphs G of order n , girth 5 and minimum degree 2 with $\gamma^{ID}(G) = \frac{3}{5}(n - 1)$.*

Proof. Consider a vertex x attached to $k \geq 2$ copies of C_5 via one of each copy's vertex (Figure 8). The set formed by three consecutive vertices of each copy of C_5 (centered in the neighbour of x) is clearly an identifying code. For the lower bound, assume that some copy contains at most two vertices of an identifying code C . Then they must be non-adjacent (otherwise some vertex is not dominated). But then at least one of these two vertices is not separated from one of its neighbours, a contradiction. Hence each copy of C_5 contains at least three vertices of C , proving the bound. \square

The following lemma is about the graph P_{11} .




Figure 8: A family of connected graphs with identifying code number $\frac{3}{5}(n - 1)$.

Lemma 26. *Let G be an identifiable graph of girth 5 containing a copy P of P_{11} as an induced subgraph, such that in P , only vertex x has neighbours out of P . Let C be an identifying code of G and $C \cap V(P) = C_P$. Then:*

- (i) $|C_P| \geq 4$;
- (ii) if $|C_P| = 4$, then x is only dominated by a vertex $y \notin V(P)$;

Proof. (i) By contradiction, assume that $|C_P| = 3$. If C_P induces a connected graph, then one can check that there are some non-dominated vertices in P . Hence, by Lemma 5(i), either C_P induces a K_2 containing x and an isolated vertex, or three isolated vertices. In both cases some vertices of P would not be separated, a contradiction.

(ii) Assume that $|C_P| = 4$ and by contradiction, that x is dominated by a vertex of C_P . If $x \notin C_P$, then C_P must form an identifying code of $P \setminus \{x\}$. Then, the bound $\gamma^{ID}(G) \geq \frac{2n}{\Delta+2}$ of Theorem 18 is tight, and by the same theorem, all vertices in C_P have degree 3 in $P \setminus \{x\}$. Hence the neighbours of x do not belong to C_P , a contradiction. Hence, $x \in C_P$.

Let $m(C_P)$ and $m(C_P, S)$ count the number of edges between vertices of C_P and edges between vertices of C_P and $S = V(P) \setminus C_P$, respectively. Let i denote the number of vertices in C_P that are not adjacent to any other vertex of C_P (note that $0 \leq i \leq 2$ since $x \in C_P$ and x is dominated by a vertex of C_P). Then, we have $m(C_P) = 4 - i - 1$ (indeed C_P must induce a forest). We also have $m(C_P, S) = 11 - 2m(C_P)$ (since $x \in C_P$ and has degree 2 in P). We get that $m(C_P, S) = 5 + 2i$. On the other hand, at most $4 - i$ vertices in S can be 1-dominated, and the other ones must be at least 2-dominated. Since $|S| = 7$, we get $m(C_P, S) \geq 4 - i + 2(7 - (4 - i)) = 10 + i$. Putting both inequalities together, we get that $i \geq 5$, a contradiction. \square

Proposition 27. *There are infinitely many connected graphs G of order n , girth 5 and minimum degree 3 with $\gamma^{ID}(G) = \frac{5}{11}(n - 1) > 0.454n$.*

Proof. Consider the graph G_{11}^k from Definition 20. An identifying code of size $5k$, formed by vertices $\{x, 2, 4, 7, 9\}$ of each copy of P_{11} , is illustrated in Figure 5(b). Now, consider an identifying code C of the graph. By Lemma 26(i), every copy of P_{11} contains at least four vertices of C . By Lemma 26(ii), for each copy of P_{11} containing *exactly* four code-vertices, then $y \in C$ and vertex x is dominated only by y . Hence there can be only one such copy, proving the lower bound. \square

We now define a cubic graph on 12 vertices with girth 5.

Definition 28. Let G_{12} be the 12-vertex graph with vertex set $\{0, 1, \dots, 11\}$ and edges given by a hamiltonian cycle $0 - 1 - 2 - \dots - 11 - 0$ and $\{0, 4\}$, $\{1, 8\}$, $\{2, 6\}$, $\{3, 10\}$, $\{5, 9\}$, and $\{7, 11\}$.

An illustration is given in Figure 9. We remark that, alternatively, G_{12} can be obtained from the Petersen graph by subdividing two edges that are at maximum distance (i.e. distance 2) from each other and joining the two new vertices by an edge. A third way is to take the Heawood graph, delete two adjacent vertices x, y and adding an edge between the two neighbours of x and an edge between the two neighbours of y .




Figure 9: The graph G_{12} with a minimum identifying code (black vertices).

Proposition 29. The graph G_{12} has $\gamma^{ID}(G_{12}) = 6 = \frac{n}{2}$.

Proof. An identifying code of size 6 is given for instance by the set $\{0, 2, 5, 6, 7, 10\}$, implying that $\gamma^{ID}(G_{12}) \leq 6$.

To prove our claim, it is sufficient to show that there is no identifying code on 5 vertices. Assume for contradiction that there is an identifying code C of G_{12} of size 5. Let I be the set of isolated vertices in C , $S = V(G_{12}) \setminus C$. Let $m(C)$ and $m(C, S)$ count the number of edges between vertices of C and the edges between C and S , respectively. By Lemma 5, at least $|S| - |C \setminus I|$ vertices from S have to be 2-dominated. Hence, there are at least $|C \setminus I| + 2(|S| - |C \setminus I|)$ edges from S to C . On the other hand, there are $3|C| - 2m(C) = 15 - 2m(C)$ edges from C to S . Hence,

$$\begin{aligned} 15 - 2m(C) = m(C, S) &\geq |C \setminus I| + 2(|S| - |C \setminus I|) \\ &= 2|S| - |C \setminus I| \\ &= 2(12 - |C|) - |C| + |I| = 9 + |I|, \end{aligned}$$

which gives

$$m(C) \leq 3 - \frac{|I|}{2}. \tag{2}$$

By Lemma 5(i), the subgraph induced by C consists either of a single component of order 5, a component of order 4 and an isolated vertex, a component of order 3 and two isolated vertices, or C is an independent set. We distinguish now between these cases.

Case a: C consists of a single component of order 5. Then $m(C) \geq 4$ and thus, by Inequality (2), $4 \leq 3$, which is a contradiction.

Case b: C consists of a component of order 4 and an isolated vertex. Again, by Inequality (2), $3 \leq m(C) \leq 2.5$, a contradiction.

Case c: C consists of a component C_c of order 3 and two isolated vertices x and y . Then C_c is a path of length 2, say uvw , and $m(C) = 2 = 3 - \frac{|I|}{2}$, giving equality in the above inequality chain. Hence, three vertices from S have exactly one neighbour in C , while the other four have exactly two neighbours in C . With $S = \{s_1, s_2, s_3, s_4, s_5, s_6, s_7\}$, let us say that s_1, s_2 , and s_3 are the vertices being dominated once and let $\{s_1, u\}, \{s_2, v\}, \{s_3, w\}$ be the edges hereby involved. Then the edges incident with v have been all assigned, while u and w can still contribute dominating one more vertex from S . However, to 2-dominate the vertices in $\{s_4, s_5, s_6, s_7\}$, necessarily two of them will be adjacent to both x and y , building a cycle of length 4, which is not allowed. Thus, this case is not possible.

Case d: $C = I = \{x_1, x_2, x_3, x_4, x_5\}$. By Lemma 5(ii), all vertices from S have to be 2-dominated by I , and hence $m(C, S) \geq 14$. But since G_{12} is cubic, each vertex of S is incident to at most one further edge. Since $|S| = 7$, at most six vertices in S can be paired, and $m(C, S) \geq 15$. On the other hand, each vertex of C has three neighbours, hence $m(C, S) \leq 15$. This implies that while one vertex from S , say s_7 , has exactly three neighbours in I , the other six vertices from S have exactly two neighbours in I . Since G_{12} is cubic, the vertices in $\{s_1, s_2, s_3, s_4, s_5, s_6\}$ are paired by a matching, say $\{s_1, s_2\}, \{s_3, s_4\}, \{s_5, s_6\}$. Consider the edge $\{s_1, s_2\}$ and its neighbours in I , say $\{x_1, x_2, x_3, x_4\}$. Going through all edges from the graph G_{12} and considering their four independent neighbours, there are only two possibilities where the corresponding independent sets can be completed to an independent set of size 5. These are the edges $\{3, 4\}$ and $\{7, 8\}$ which give each two possible independent sets $\{0, 2, 5, 7, 10\}, \{0, 2, 5, 8, 10\}$ and $\{1, 3, 6, 9, 11\}, \{1, 4, 6, 9, 11\}$. Hence, $C = I$ has to be one of these sets. However, it is easy to check that none of them is an identifying code.

Hence, G_{12} has no identifying code of size 5 and $\gamma^{\text{ID}}(G_{12}) = 6$. □

4 Conclusion

We proved the two tight upper bounds $\gamma^{\text{LD}}(G) \leq \frac{n}{2}$ and $\gamma^{\text{ID}}(G) \leq \frac{5}{7}n$ for graphs G of girth at least 5 and minimum degree at least 2, as well as improved bounds for cubic graphs. While the first bound is asymptotically tight for large values of n , we do not know whether this holds for the latter one.

For minimum degree at least 3, either our bounds are not tight, or we have not found the graphs with highest value of parameters γ^{LD} and γ^{ID} . In particular, the question whether for every graph G of girth at least 5 and minimum degree at least 3, we have

$\gamma^{\text{ID}}(G) \leq \frac{n}{2}$ seems intriguing. By Proposition 29 this would be tight for the graph G_{12} . Though we have tried to get better bounds when $\delta \geq 3$, it seems that our technique is not powerful enough for such an improvement (at least without any new idea).

To conclude, we remark that another interesting question would be to conduct a similar study for the *open location-domination number* and the *locating-total domination number*, related concepts introduced in [23] and [12], respectively.

Acknowledgements. We thank the anonymous referee for carefully reading the paper, therefore helping to improve its presentation.

References

- [1] N. Alon and J. H. Spencer. *The probabilistic method*, 3rd edition, Wiley-Interscience, 2008.
- [2] N. Alon. Transversal numbers of uniform hypergraphs, *Graphs and Combinatorics* 6(1):1–4, 1990.
- [3] L. Babai. On the complexity of canonical labeling of strongly regular graphs. *SIAM Journal on Computing* 9(1):212–216, 1980.
- [4] N. Bertrand, I. Charon, O. Hudry, A. Lobstein. Identifying and locating-dominating codes on chains and cycles. *European Journal of Combinatorics* 25(7):969–987, 2004.
- [5] B. Bollobás and A. D. Scott. On separating systems. *European Journal of Combinatorics* 28:1068–1071, 2007.
- [6] J. A. Bondy. Induced subsets. *Journal of Combinatorial Theory, Series B* 12(2):201–202, 1972.
- [7] K. M. J. De Bontridder, B. V. Halldórsson, M. M. Halldórsson, C. A. J. Hurkens, J. K. Lenstra, R. Ravi, L. Stougie. Approximation algorithms for the test cover problem. *Mathematical Programming Series B* 98:477–491, 2003.
- [8] E. Charbit, I. Charon, G. Cohen, O. Hudry, A. Lobstein. Discriminating codes in bipartite graphs: bounds, extremal cardinalities, complexity. *Advances in Mathematics of Communications* 2(4):403–420, 2008.
- [9] F. Foucaud. *Combinatorial and algorithmic aspects of identifying codes in graphs*. PhD thesis, Université Bordeaux 1, France, December 2012.
<http://tel.archives-ouvertes.fr/tel-00766138>
- [10] F. Foucaud and G. Perarnau. Bounds for identifying codes in terms of degree parameters. *Electronic Journal of Combinatorics* 19:#P32, 2012.
- [11] D. Garijo, A. González, A. Márquez. The difference between the metric dimension and the determining number of a graph. *Applied Mathematics and Computation* 249:487–501, 2014.
- [12] T. W. Haynes, M. A. Henning, J. Howard. Locating and total dominating sets in trees. *Discrete Applied Mathematics* 154:1293–1300, 2006.

- [13] M. G. Karpovsky, K. Chakrabarty, L. B. Levitin. On a new class of codes for identifying vertices in graphs. *IEEE Transactions on Information Theory* 44:599–611, 1998.
- [14] J. H. Kim, O. Pikhurko, J. Spencer, O. Verbitsky. How complex are random graphs in First Order logic? *Random Structures and Algorithms* 26(1-2):119–145, 2005.
- [15] A. V. Kostochka and B. Y. Stodolsky. An upper bound on the domination number of n -vertex connected cubic graphs. *Discrete Mathematics* 309:1142–1162, 2009.
- [16] M. Laifenfeld, A. Trachtenberg, R. Cohen, D. Starobinski. Joint monitoring and routing in wireless sensor networks using robust identifying codes. *Proceedings of IEEE Broadnets 2007*, pages 197–206, September 2007.
- [17] A. Lobstein. Watching systems, identifying, locating-dominating and discriminating codes in graphs: a bibliography.
<http://www.infres.enst.fr/~lobstein/debutBIBidetlocdom.pdf>
- [18] W. McCuaig and B. Shepherd. Domination in graphs with minimum degree two. *Journal of Graph Theory* 13(6):749–762, 1989.
- [19] B. M. E. Moret and H. D. Shapiro. On minimizing a set of tests. *SIAM Journal on Scientific and Statistical Computation* 6(4):983–1003, 1985.
- [20] O. Ore. Theory of graphs. American Mathematical Society Colloquium Publications Vol. XXXVIII, American Mathematical Society, Providence, R.I., 1962.
- [21] B. Reed. Paths, stars and the number three. *Combinatorics, Probabilities and Computing* 5(3):277–295, 1996.
- [22] A. Rényi. On random generating elements of a finite Boolean algebra. *Acta Scientiarum Mathematicarum Szeged* 22:75–81, 1961.
- [23] S. J. Seo and P. J. Slater. Open neighborhood locating-dominating sets. *The Australasian Journal of Combinatorics* 46:109–120, 2010.
- [24] P. J. Slater. Domination and location in acyclic graphs. *Networks* 17(1):55–64, 1987.
- [25] P. J. Slater. Dominating and reference sets in a graph. *Journal of Mathematical and Physical Sciences* 22(4):445–455, 1988.
- [26] P. J. Slater. Locating dominating sets and locating-dominating sets. Graph Theory, Combinatorics, and Applications: Proceedings of the 7th Quadrennial International Conference on the Theory and Applications of Graphs 2:1073–1079, Wiley, 1995.
- [27] R. Ungrangsi, A. Trachtenberg, D. Starobinski. An implementation of indoor location detection systems based on identifying codes. *Proceedings of Intelligence in Communication Systems, INTELLCOMM 2004, Lecture Notes in Computer Science* 3283:175–189, 2004.
- [28] H. Xing, L. Sun, X. Chen. Domination in graphs of minimum degree five. *Graphs and Combinatorics* 22(1):127–143, 2006.