

# Hitting minors, subdivisions, and immersions in tournaments

Jean-Florent Raymond\*

## Abstract

The Erdős–Pósa property relates parameters of covering and packing of combinatorial structures and has been mostly studied in the setting of undirected graphs. In this note, we use results of Chudnovsky, Fradkin, Kim, and Seymour to show that, for every directed graph  $H$  (resp. strongly-connected directed graph  $H$ ), the class of directed graphs that contain  $H$  as a strong minor (resp. butterfly minor, topological minor) has the vertex-Erdős–Pósa property in the class of tournaments. We also prove that if  $H$  is a strongly-connected directed graph, the class of directed graphs containing  $H$  as an immersion has the edge-Erdős–Pósa property in the class of tournaments. Our results are orthogonal to the recent results of Amiri et al. [arXiv:1603.02504, March 2016] in the sense that we restrict the class of “host graphs”, whereas they restrict the class of “guest graphs”.

**Keywords:** directed Erdős–Pósa property, packing and covering, topological minors, immersions, tournaments.

## 1 Introduction

We are concerned in this note with the Erdős–Pósa property in the setting of directed graphs. This property, which has mostly been studied on undirected graphs, is originated from the following classic result by Erdős and Pósa: there is a function  $f: \mathbb{N} \rightarrow \mathbb{N}$ , such that, for every (undirected) graph  $G$  and every positive integer  $k$ , one of the following holds: (a)  $G$  contains  $k$  vertex-disjoint cycles; or (b) there is a set  $X \subseteq V(G)$  with  $|X| \leq f(k)$  and such that  $G \setminus X$  has no cycle [8]. This theorem expresses a duality between a parameter of *packing*, the maximum number of vertex-disjoint cycles in a graph, and a parameter of *covering*, the minimum number of vertices that intersects all cycles. This initiated a research line aimed at providing conditions for this property to hold, for various combinatorial objects. Formally, we say that a class  $\mathcal{H}$  has the *Erdős–Pósa property* if there is a function  $f: \mathbb{N} \mapsto \mathbb{N}$  such that, for every positive integer  $k$  and every graph  $G$  (referred to as the *host graph*) one of the following holds:

- $G$  has  $k$  vertex-disjoint subgraphs that are isomorphic to members of  $\mathcal{H}$ ; or
- there is a set  $X \subseteq V(G)$  with  $|X| \leq f(k)$  and such that  $G \setminus X$  has no subgraph isomorphic to a member of  $\mathcal{H}$ .

The Erdős–Pósa Theorem states that the class of cycles has this property. One of the most general extensions of the Erdős–Pósa Theorem is certainly the following byproduct of the Graph Minors series: Robertson and Seymour proved that the class of graphs that contain  $H$  as a minor have the Erdős–Pósa property iff  $H$  is planar [23].

---

\*LIRMM, Montpellier, France and Institute of Computer Science, University of Warsaw, Poland. This research has been supported by the grant PRELUDIUM 2013/11/N/ST6/02706 of the Polish National Science Center (NCN). Email: [jean-florent.raymond@mimuw.edu.pl](mailto:jean-florent.raymond@mimuw.edu.pl)

On the other hand, some classes fail to have the Erdős–Pósa property, as odd cycles [6]. When this happens, one can investigate on particular classes of host graphs. In this direction, Reed [19] proved that odd cycles have the Erdős–Pósa property in planar graphs.

A natural variant of the Erdős–Pósa property is to change, in the definition, vertex-disjoint subgraphs for edge-disjoint ones and sets of vertices for sets of edges. It appears that the Erdős–Pósa Theorem also holds in this setting [7]. Other results have been obtained about this variant, less than on the vertex variant, though. At this point we have to stress that, if the vertex and edge variants of the Erdős–Pósa property have close definitions, one cannot in general deduce one from the other. We refer the reader to surveys [18, 20] for more details about the Erdős–Pósa property.

In the setting of directed graphs however, few results are known. Until recently, the largest class of such graphs that has been studied under the prism of the Erdős–Pósa property was the class of directed cycles [11, 13, 21, 22, 24]. It is worth noting that, besides its combinatorial interest, the Erdős–Pósa property in directed graphs found applications in bioinformatics and in the study of Boolean networks [3, 4]. We consider here finite directed graphs (*digraphs*) that may have multiple arcs, but not loops and we respectively denote by  $V(G)$  and  $E(G)$  the set of vertices and the multiset of arcs of a digraph  $G$ . A digraph  $G$  is said to be *strongly-connected* if it has at least one vertex and, for every  $u, v \in V(G)$ , there is a directed path from  $u$  to  $v$ . The most general result about the Erdős–Pósa property in digraphs is certainly the following directed counterpart of the aforementioned results of Robertson and Seymour.

**Theorem 1** ([2]). *Let  $H$  be a strongly-connected digraph that is a butterfly minor (resp. topological minor) of a cylindrical grid.<sup>1</sup> There is a function  $f: \mathbb{N} \rightarrow \mathbb{N}$ , such that for every digraph  $G$  and every positive integer  $k$ , one of the following holds:*

- $G$  has  $k$  vertex-disjoint subdigraphs, each having  $H$  as a butterfly minor (resp. topological minor); or
- there is a set  $X \subseteq V(G)$  with  $|X| \leq f(k)$  such that  $G \setminus X$  does not have  $H$  as a butterfly minor (resp. topological minor).

On the other hand, the authors of [2] proved that **Theorem 1** does not hold for the strongly-connected digraphs  $H$  that do not satisfy the conditions of its statement. It seems therefore natural to ask under what restrictions on the host digraphs the above result could be true for every strongly-connected digraph, in the same spirit as the aforementioned result of Reed.

The purpose of this note is twofold: obtaining new Erdős–Pósa type results on directed graphs and providing evidence that techniques analogues to those used in the undirected case may be adapted to the directed setting. In particular, we describe conditions on the class of host digraphs so that **Theorem 1** holds for every strongly-connected digraph  $H$ . Before we formally state our results, let us introduce some terminology.

Several directed counterparts of the notion of minor have been introduced in the literature. An arc  $(u, v)$  of a digraph is said to be *contractible* if either it is the only arc with head  $v$ , or it is the only arc with tail  $u$ . Following [14] and [16], we say that a digraph  $H$  is a *butterfly minor* (resp. *strong minor*) of a digraph  $G$  if a digraph isomorphic to  $H$  can be obtained from a subdigraph of  $G$  by contracting contractible arcs (resp. contracting strongly-connected subdigraphs to single vertices). Notice that these notions are incomparable. A motivation for these definitions is that taking (butterfly or strong) minors does not create directed cycles. Unlike minors, immersions and topological minors are concepts that are easily extended to the setting of directed graphs as they can be defined in terms of paths. We say that a digraph  $H$  is a *topological minor* of a digraph  $G$  if there is a

---

<sup>1</sup>The notions of butterfly minor and topological minor will be defined in a forthcoming paragraph. We refer the reader to [2] for a definition of the cylindrical directed grid.

subdigraph of  $G$  that can be obtained from a digraph isomorphic to  $H$  by replacing arcs by directed paths (in the same direction) that do not share internal vertices. If we allow these paths to share internal vertices but not arcs, then we say that  $H$  is an *immersion* of  $G$ . Observe that every topological minor is a butterfly minor. However, as often with the Erdős–Pósa property, this does not allow us in general to deduce an Erdős–Pósa-type result about the one relation from a result about the other one.

Our results hold on superclasses of the extensively studied class of *tournaments*, that are all orientations of undirected complete graphs. For  $s \in \mathbb{N}$ , a digraph is *s-semicomplete* if, for every vertex  $v$ , there are at most  $s$  vertices that are not connected to  $v$  by an arc (in either direction). A *semicomplete* digraph is a 0-semicomplete digraph. These classes generalize the class of tournaments. Our contributions are the following two theorems.

**Theorem 2.** *For every finite family  $\mathcal{H}$  of strongly-connected digraphs and every  $s \in \mathbb{N}$ , there is a function  $f: \mathbb{N} \rightarrow \mathbb{N}$  such that for every  $s$ -semicomplete digraph  $G$  and every positive integer  $k$ , one of the following holds:*

- $G$  has  $k$  vertex-disjoint subdigraphs, each having a digraph of  $\mathcal{H}$  as a strong minor (resp. butterfly minor, topological minor); or
- there is a set  $X \subseteq V(G)$  with  $|X| \leq f(k)$  such that  $G \setminus X$  does not have a digraph of  $\mathcal{H}$  as a strong minor (resp. butterfly minor, topological minor).

When considering the strong minor relation, the above result also holds when digraphs in  $\mathcal{H}$  are not strongly-connected (see [Lemma 4](#)).

**Theorem 3.** *For every family  $\mathcal{H}$  of strongly-connected digraphs on at least two vertices, there is a function  $f: \mathbb{N} \rightarrow \mathbb{N}$  such that for every semicomplete  $G$  and every positive integer  $k$ , one of the following holds:*

- $G$  has  $k$  arc-disjoint subdigraphs, each having a digraph of  $\mathcal{H}$  as an immersion; or
- there is a set  $X \subseteq E(G)$  with  $|X| \leq f(k)$  such that  $G \setminus X$  does not have a digraph of  $\mathcal{H}$  as an immersion.

These theorems deal with the two variants of the Erdős–Pósa property: the first one is related to vertex-disjoint subdigraphs and sets of vertices (vertex version), whereas the second one is concerned with arc-disjoint subdigraphs and sets of arcs (arc version). In [Theorem 3](#), the requirement on the size of the graphs of  $\mathcal{H}$  is necessary as we cannot cover an arcless subdigraph (as the one-vertex digraph) with arcs. Our proofs rely on exclusion results for the parameters of cutwidth and pathwidth, that are stated in the sections where they are used.

The techniques that we use are originated from the undirected setting, where they have been repeatedly applied (see for instance [\[23, \(8.8\)\]](#) and [\[9, Lemma 2.3\]](#)). They deal with structural decompositions like tree decompositions or tree-cut decompositions and their associated widths and can be informally described as follows. If the considered host graph has large width, then, using a structural result, we can immediately conclude that it contains several disjoint subgraphs of the desired type. Otherwise, the graph admits a structural decomposition of small width, that can be used to find a small set of vertices/edges covering all such subgraphs (see [\[18, Theorem 3.1\]](#) for an unified presentation in undirected graphs). Similar ideas have been used in the context of directed graphs in the proof of [Theorem 1](#). With this note, we provide more examples of cases where the techniques used in the undirected setting appear useful when dealing with digraphs.

## 2 Hitting minors and subdivisions

The definitions of (butterfly and strong) minors and topological minors have been given in the introduction. The structural decompositions that we use in this section are path

decompositions. Formally, a *path-decomposition* of a digraph  $G$  is a sequence  $(X_1, \dots, X_r)$  of subsets of  $V(G)$  satisfying the following properties:

- (i)  $V(G) = \bigcup_{i=1}^r X_i$
- (ii) for every arc  $(u, v) \in E(G)$ , there are integers  $i \in \{1, \dots, r\}$  and  $j \in \{1, \dots, i\}$  such that  $u \in X_i$  and  $v \in X_j$ ;
- (iii) for every  $i, j \in \{1, \dots, r\}$ , if a vertex  $u \in V(G)$  belongs to  $X_i$  and  $X_j$ , then it also belongs to  $X_k$  for every  $k \in \{i, \dots, j\}$ .

The sets  $\{X_i\}_{i \in \{1, \dots, r\}}$  are called *bags* of the path-decomposition. Intuitively, item (ii) asks that every arc of  $G$  either have its endpoints in some bag, or is oriented “backwards”. The *width* of the above path-decomposition is defined as  $\max_{i \in \{1, \dots, r\}} |X_i| - 1$ . The *pathwidth* of  $G$  is the minimum width over all path-decompositions of  $G$ . The following properties of pathwidth are crucial in our proofs.

**Theorem 4** ([15, Theorem 2.2.7], [10, (1.1)], [16, (1.4)]). *For every digraph  $H$ , there is a positive integer  $w$  such that every semicomplete  $G$  that has pathwidth more than  $w$  contains  $H$  as a strong minor, butterfly minor and topological minor.*

**Theorem 5** ([17, Theorem 2]). *For every  $s, w \in \mathbb{N}$ , there is a positive integer  $w'$  such that every  $s$ -semicomplete digraph with pathwidth at least  $w'$  has a subdigraph that is semicomplete and is of pathwidth at least  $w$ .*

**Corollary 1.** *For every  $s \in \mathbb{N}$  and every digraph  $H$ , there is a constant  $\zeta_{s,H}$  such that every  $s$ -semicomplete digraph that has pathwidth at least  $\zeta_{s,H}$  contains  $H$  as a strong minor, butterfly minor and topological minor.*

A classic result states that if a collection of subpaths of a path does not contain more than  $k$  vertex-disjoint elements, then there is a set of  $k - 1$  vertices meeting all the subpaths [12]. We use here the following generalization of the above statement.

**Lemma 1** ([1]). *Let  $P$  be a path (undirected) and let  $\mathcal{P}$  be a collection of subgraphs of  $P$  that does not contain  $k + 1$  pairwise vertex-disjoint members. Then there is a set of  $2p^2k$  vertices of  $P$  meeting every element of  $\mathcal{P}$ , where  $p$  is the maximal number of connected components of a graph in  $\mathcal{P}$ .*

A *strongly-connected component* of a digraph is a maximal subdigraph that is strongly-connected. If a subdigraph of a digraph  $G$  is isomorphic to some member of a digraph class  $\mathcal{H}$ , we call it an  $\mathcal{H}$ -subdigraph of  $G$ .

**Lemma 2.** *Let  $\mathcal{H}$  be a class of digraphs with at most  $p$  strongly-connected components. For every digraph  $G$  and a positive integer  $k$ , one of the following holds: (a)  $G$  contains  $k$  pairwise vertex-disjoint  $\mathcal{H}$ -subdigraphs; or (b) there is a set  $X \subseteq V(G)$  with  $|X| \leq 2p^2(k - 1)(\mathbf{pw}(G) + 1)$  such that  $G \setminus X$  has no  $\mathcal{H}$ -subdigraph.*

*Proof.* We proceed by induction on  $k \in \mathbb{N}$ . The base case  $k = 1$  is trivial. Let us prove the statement of the lemma for  $k > 1$  assuming that it holds for all lower values of  $k$  (induction step). For this we consider a digraph  $G$  such that (a) does not hold. Let  $(X_1, \dots, X_l)$  be a path decomposition of  $G$  of minimum width. Let  $P$  be the undirected path on vertices  $v_1, \dots, v_l$ , in this order. For every subdigraph  $F$  of  $G$ , we set:

$$A_F = \{i \in \{1, \dots, l\}, V(F) \cap X_i \neq \emptyset\} \quad \text{and} \\ P_F = P[\{v_i, i \in A_F\}].$$

In other words,  $A_F$  is the set of indices of the bags met by  $F$  and  $P_F$  is the subgraph of  $P$  induced by the vertices with these indices. For every  $\mathcal{H}$ -subdigraph  $H$  of  $G$ , we consider

the subgraph  $P_H$  of  $P$ . Let us denote by  $\mathcal{P}$  the class of all such graphs. Notice that for every pair  $F, F'$  of subdigraphs of  $G$ , if  $P_F$  and  $P_{F'}$  are vertex-disjoint, then so are  $F$  and  $F'$ . Using our initial assumption on  $G$ , we deduce that  $\mathcal{P}$  does not contain  $k$  pairwise vertex-disjoint members. Besides, if  $F$  is strongly-connected, then  $P_F$  is connected. Moreover, if  $F$  has at most  $p$  strongly-connected components, then  $P_F$  has at most  $p$  connected components. Hence, every member of  $\mathcal{P}$  has at most  $p$  connected components.

By the virtue of **Lemma 1**, there is a set  $Q$  of  $2p^2(k-1)$  vertices of  $P$  such that  $P \setminus Q$  does not contain a subgraph of  $\mathcal{P}$ . Let  $X = \bigcup_{i \in \{j \in \{1, \dots, l\}, v_j \in Q\}} X_i$ . Let us show that  $X$  satisfies the requirements of **(b)**. By contradiction, we assume that  $G \setminus X$  has an  $\mathcal{H}$ -subdigraph  $H$ . Then  $P_H \in \mathcal{P}$ . Let  $v_i$  be a vertex of  $V(P_H) \cap Q$ , which, by definition of  $Q$ , is not empty. Then  $X_i \subseteq X$  and  $V(H) \cap X_i \neq \emptyset$ . This contradicts the fact that  $H$  is a subdigraph of  $G \setminus X$ . Consequently,  $X$  is as required. As it is the union of  $2p^2(k-1)$  bags of an optimal path decomposition of  $G$ , we have  $|X| \leq 2p^2(k-1)(\mathbf{pw}(G) + 1)$ . This concludes the proof.  $\square$

We would like to mention that a weaker form of **Lemma 2** where  $\mathcal{H}$  consists of digraphs whose connected components are strongly-connected can be obtained by adapting the ideas used in [23, (8.8)], with a dependency in  $p$  that is linear instead of quadratic.

**Lemma 3.** *Let  $G$  be a digraph and let  $H$  be the digraph obtained by contracting one strongly-connected subdigraph  $S$  of  $G$  to one single vertex  $v_S$ . Then  $H$  and  $G$  have the same number of strongly-connected components.*

*Proof.* Let  $f$  be the map such that, for every  $C \subseteq V(H)$  that induces a strongly-connected component,

$$f(C) = \begin{cases} C & \text{if } v_S \notin C \\ (C \setminus \{v_S\}) \cup S & \text{otherwise} \end{cases}$$

Let  $C$  be a subset of  $V(H)$  that induces a strongly-connected component and let us show that  $f(C)$  induces a strongly-connected subdigraph of  $G$ . For this, we show that, for any  $x, y \in f(C)$ , there is a directed path from  $x$  to  $y$ . If  $x, y \in S$ , this is true since in this case,  $S \subseteq f(C)$  and  $G[S]$  is strongly-connected. If none of  $x, y$  belongs to  $S$ , they are both vertices of  $C$  as well. Let  $v_0 \dots v_l$  be a directed path from  $x = v_0$  to  $y = v_l$  in  $H$ . If  $v_S$  does not belong to this path, then this is a path of  $G$  as well and we are done. Otherwise, let  $i$  be such that  $v_S = v_i$ . By definition of  $H$ , there are arcs  $(v_{i-1}, u)$  and  $(u', v_{i+1})$  in  $G$ , for some  $u, u' \in S$ . As  $G[S]$  is strongly-connected, it contains a directed path  $Q$  from  $u$  to  $u'$ . Therefore, concatenating  $v_0 \dots v_{i-1}u$ ,  $Q$ , and  $u'v_{i+1} \dots v_l$  yields a path from  $x$  to  $y$ . The case where one of  $x, y$  belong to  $S$  is similar. Consequently,  $f(C)$  induces a strongly-connected subdigraph in  $G$ .

Let us now show that  $f(C)$  is a strongly-connected component. By contradiction, let us assume that there is in  $G$  a directed walk  $u_0 \dots u_l$  with  $l > 1$  such that  $\{u_0 \dots u_l\} \cap f(C) = \{u_0, u_l\}$ . If  $S \cap (f(C) \cup \{u_0 \dots u_l\}) = \emptyset$  then  $C$  does not induce a strongly-connected component of  $H$ , a contradiction. If  $S \cap f(C) = \emptyset$ , then  $f(C) = C$  and then  $G$  has a path (that is a subpath of  $u_0 \dots u_l$ ) from a vertex of  $f(C)$  to one of  $S$  and vice-versa. Therefore, there is in  $H$  a path from a vertex of  $C$  to  $v_S$  and vice-versa, which contradicts the definition of  $C$ . Therefore  $S$  intersects  $f(C)$ : by definition of  $f$  we have  $S \subseteq f(C)$  and  $v_S \in C$ . We deduce that  $P = u_0 \dots u_l$  (or  $P = v_S, u_1 \dots u_l$ , resp.  $P = u_0 \dots u_{l-1}, v_S$  if  $u_0 \in S$ , resp.  $u_l \in S$ ) is an oriented walk of  $H$  on at least 3 vertices with  $|C \cap V(P)| < 3$ , which is not possible since  $C$  induces a strongly-connected component of  $H$ . We deduce that  $f(C)$  is a strongly-connected component of  $G$ .

The function  $f$  is clearly injective. Let us show that it is surjective. Let now  $C \subseteq V(G)$  be a strongly-connected component of  $G$ . If  $C$  contains a vertex of  $S$ , then  $S \subseteq C$  as  $C$  is a maximal strongly-connected subdigraph and  $S$  is strongly-connected. In this case observe that  $f(C \setminus S) \cup \{v_S\} = C$ . Otherwise,  $C \cap S = \emptyset$  and  $f(C) = C$ . Strongly-connected components of  $G$  are in bijection with those of  $H$ , hence they are equally many.  $\square$

**Corollary 2.** *Let  $H$  be a digraph and let  $G$  be a subdigraph-minimal digraph containing  $H$  as a strong minor. Then  $H$  and  $G$  have the same number of strongly-connected components.*

**Lemma 4.** *For every family  $\mathcal{H}$  of digraphs with bounded number of strongly-connected components and every  $s \in \mathbb{N}$ , there is a function  $f: \mathbb{N} \rightarrow \mathbb{N}$  such that, for every  $s$ -semicomplete digraph  $G$  and every positive integer  $k$ , one of the following holds: (a)  $G$  contains  $k$  vertex-disjoint subdigraphs, each having a digraph of  $\mathcal{H}$  as a strong minor; or (b) there is a set  $X \subseteq V(G)$  with  $|X| \leq f(k)$  such that  $G \setminus X$  contains no digraph of  $\mathcal{H}$  as a strong minor.*

*Proof.* Let us assume that (a) does not hold (otherwise we are done). Let  $H \in \mathcal{H}$ . According to [Corollary 1](#), we have  $\mathbf{pw}(G) < \zeta_{s,k,H}$ . Let  $p$  denote the maximum number of strongly-connected components of a digraph in  $\mathcal{H}$  and let  $\hat{\mathcal{H}}$  be the class of all subdigraph-minimal digraphs containing a digraph of  $\mathcal{H}$  as a strong minor. Observe that  $G$  has a digraph of  $\mathcal{H}$  as a strong minor iff it has a subgraph isomorphic to a digraph in  $\hat{\mathcal{H}}$ . Also, according to [Corollary 2](#), the digraphs in  $\hat{\mathcal{H}}$  have at most  $p$  strongly-connected components. We can now apply [Lemma 2](#) and obtain a set  $X$  of at most  $2p^2(k-1)\zeta_{s,k,H}$  vertices such that  $G \setminus X$  contains no digraph of  $\mathcal{H}$  as a strong minor, that is, item (b). This concludes the proof.  $\square$

In general, digraphs containing a digraph  $H$  as a butterfly minor (resp. topological minor) may have more strongly-connected components than  $H$ . Therefore we focus on strongly-connected digraphs where the following result plays the role of [Lemma 3](#).

**Lemma 5.** *Let  $G$  be a strongly-connected digraph and let  $H$  be the digraph obtained by contracting a contractible arc  $(s, t)$  of  $G$ . Then  $H$  is strongly-connected.*

*Proof.* Notice that case where  $H$  is a single vertex is not possible as this digraph has no arc. Towards a contradiction, let us assume that there are two vertices  $x, y \in V(H)$  such that there is a directed path  $v_1 \dots v_l$  from  $x = v_1$  to  $y = v_l$  in  $G$  but not in  $H$ . As  $G$  and  $H$  differ only by the contraction of  $(s, t)$ , there are distinct  $i, j \in \{1, \dots, l\}$  such that  $s = v_i$  and  $t = v_j$ . If  $i < j$ , then  $v_1 \dots v_i v_j \dots v_l$  is a directed path from  $x$  to  $y$  in  $H$ , a contradiction. Let us now assume that  $i > j$ . Observe that since  $G$  is strongly-connected, there are arcs  $(v_0, v_1)$  and  $(v_l, v_{l+1})$  (for some vertices  $v_0, v_{l+1}$  that may belong to the path we consider). Now,  $v_i$  is the tail of the two arcs  $(v_i, v_j)$  and  $(v_i, v_{i+1})$  and  $v_j$  is the head of the two arcs  $(v_i, v_j)$  and  $(v_{j-1}, v_j)$ , which contradicts the contractibility of  $(s, t)$ . Therefore,  $H$  is strongly-connected.  $\square$

**Corollary 3.** *Let  $H$  be a digraph whose connected components are strongly-connected and let  $G$  be a subdigraph-minimal digraph containing  $H$  as a butterfly minor (resp. topological minor). Then  $H$  and  $G$  have the same number of strongly-connected components.*

**Lemma 6.** *For every finite family  $\mathcal{H}$  of digraphs whose connected components are strongly-connected and every  $s \in \mathbb{N}$ , there is a function  $f: \mathbb{N} \rightarrow \mathbb{N}$  such that, for every  $s$ -semicomplete digraph  $G$  and every positive integer  $k$ , one of the following holds: (a)  $G$  contains  $k$  vertex-disjoint subdigraphs, each having a digraph of  $\mathcal{H}$  as a butterfly minor (resp. topological minor); or (b) there is a set  $X \subseteq V(G)$  with  $|X| \leq f(k)$  such that  $G \setminus X$  contains no digraph of  $\mathcal{H}$  as a butterfly minor (resp. topological minor).*

*Proof.* This proof is similar to that of [Lemma 4](#). Again, we can assume that (a) does not hold and deduce  $\mathbf{pw}(G) < \zeta_{s,k,H}$  from [Corollary 1](#), for some  $H \in \mathcal{H}$ . We denote by  $p$  the maximum number of connected components of a digraph in  $\mathcal{H}$  and by  $\hat{\mathcal{H}}$  the class of all subdigraph-minimal digraphs containing a digraph of  $\mathcal{H}$  as a butterfly minor (resp. strong minor). The digraphs in  $\hat{\mathcal{H}}$  have at most  $p$  strongly-connected components, according to [Corollary 3](#). We now apply [Lemma 2](#) and obtain a set  $X$  of at most  $2p^2(k-1)\zeta_{s,k,H}$  vertices satisfying item (b).  $\square$

[Theorem 2](#) is a consequence of [Lemma 4](#) and [Lemma 6](#).

### 3 Hitting immersions

The notion of immersion has been defined in the introduction. For every two subsets  $X, Y \subseteq V(G)$ , we denote by  $E_G(X, Y)$  the set of arcs of  $G$  of the form  $(x, y)$  with  $x \in X$  and  $y \in Y$ . Recall that we call  $\mathcal{H}$ -subdigraph of  $G$  every subdigraph of a digraph  $G$  that is isomorphic to some digraph in the class of digraphs  $\mathcal{H}$ . The parameter that plays a major role in this section is the cutwidth. If  $G$  is a digraph on  $n$  vertices, the *width* of an ordering  $v_1 \dots, v_n$  of its vertices is defined as

$$\max_{i \in \{2, \dots, n\}} |E_G(\{v_1, \dots, v_{i-1}\}, \{v_i, \dots, v_n\})|.$$

The *cutwidth* of  $G$ , that we write  $\mathbf{ctw}(G)$ , is the minimum cutwidth over all orderings  $V(G)$ . Intuitively, a digraph that has small cutwidth has an ordering where the number of "left-to-right" arcs is small. The following result plays a similar role as [Theorem 4](#) in the previous section.

**Theorem 6** ([5, (1.2)]). *For every digraph  $H$ , there is a positive integer  $\eta_H$  such that every semicomplete  $G$  that has cutwidth at least  $\eta_H$  contains  $H$  as an immersion.*

*Proof of Theorem 3.* Let  $H \in \mathcal{H}$  and let  $\hat{\mathcal{H}}$  the class of all subdigraph-minimal digraphs containing a digraph of  $\mathcal{H}$  as an immersion and observe that these digraphs are strongly-connected. Again,  $G$  has a digraph of  $\mathcal{H}$  as an immersion iff it has an  $\hat{\mathcal{H}}$ -subdigraph.

According to [Theorem 6](#), we are done if  $\mathbf{ctw}(G) \geq \eta_{k \cdot H}$ . Therefore we now consider digraphs of cutwidth at most  $\eta_{k \cdot H}$ .

We will prove the statement on digraphs of cutwidth at most  $t$  by induction on  $k$  with  $f: k \mapsto k \cdot t$ . The case  $k = 0$  is trivial, therefore we assume  $k > 0$  and that the result holds for every every  $k' < k$ . We also assume that  $G$  does not contain  $k$  arc-disjoint  $\hat{\mathcal{H}}$ -subdigraphs, otherwise we are done. Let  $i \in \mathbb{N}$  be the minimum integer such that  $G[v_1, \dots, v_i]$  has an  $\hat{\mathcal{H}}$ -subdigraph. Notice that  $i > 1$  as we assume that the digraphs in  $\mathcal{H}$  has at least two vertices. We set  $Y = E_G(\{v_1, \dots, v_{i-1}\}, \{v_i, \dots, v_n\})$ . Notice that  $|Y| \leq t$ . As the digraphs in  $\hat{\mathcal{H}}$  are strongly-connected, any  $\hat{\mathcal{H}}$ -subdigraph of  $G \setminus Y$  belongs to exactly one of  $\{v_1, \dots, v_i\}, \{v_{i+1}, \dots, v_n\}$ . By definition of  $i$ , every such subdigraph belongs to  $\{v_{i+1}, \dots, v_n\}$ . Notice that every subdigraph of  $\{v_{i+1}, \dots, v_n\}$  is arc-disjoint with  $J$ . Therefore  $G[\{v_{i+1}, \dots, v_n\}]$  does not contains  $k - 1$  arc-disjoint  $\hat{\mathcal{H}}$ -subdigraphs. It is clear that this subdigraph has cutwidth at most  $t$ . By induction hypothesis, there is a set  $Y' \subseteq E(G[\{v_{i+1}, \dots, v_n\}])$  such that  $G[\{v_{i+1}, \dots, v_n\}] \setminus Y'$  has no  $\hat{\mathcal{H}}$ -subdigraph and  $|Y'| \leq (k - 1) \cdot t$ . We deduce that  $G \setminus (Y \cup Y')$  has no  $\hat{\mathcal{H}}$ -subdigraph and that  $|Y \cup Y'| \leq k \cdot t$ , as required. This concludes the induction. We saw above that we only need to consider digraphs of cutwidth at most  $\eta_{k \cdot H}$  and we just proved that in this case there is a suitable set of arcs of at most  $k\eta_{k \cdot H}$ . This concludes the proof.  $\square$

### 4 Discussion

In this note we obtained new Erdős–Pósa type results about classes defined by the relations of strong minors, butterfly minors, topological minors and immersions. The restriction of the host class to tournaments (or slightly larger classes) allowed us to obtain results for every strongly-connected pattern  $H$ . In particular, we provided conditions on the host class where [Theorem 1](#) holds for every strongly-connected digraph  $H$ , which is not the case in general. Our proofs support the claim that techniques analogues to those used in the undirected case may be adapted to the directed setting. Let us now highlight two directions for future research.

**Optimization of the gap.** The bounds on the function  $f$  in our results (gap of the Erdős–Pósa property) depend on the exclusion bounds of [Theorem 4](#) and [Theorem 6](#). Therefore, any improvement of these bounds yields an improvement of  $f$ . The upper bound on  $\eta_H$  of [Theorem 6](#) that can be obtained from the proof (in [\[5\]](#)) is  $72 \cdot 2^{2h(h+2)} + 8 \cdot 2^{h(h+2)}$ , where  $h = |V(H)| + 2|E(H)|$ . As a consequence, we have  $f(k) = 2^{\mathcal{O}(k^2h^2)}$  in [Theorem 3](#). It would be interesting to know whether a gap that is polynomial in  $k$  can be obtained. The same question can be asked for [Theorem 2](#), however the upper bound in [Theorem 4](#) that we can compute from the proof in [\[10\]](#) is large (triply exponential).

**Generalization.** The results presented in this note were related to (generalizations of) semicomplete digraphs. One direction for future research would be to extend them to wider classes of hosts. On the other hand, in [Theorem 3](#), we require the guest digraph to be strongly connected when dealing with butterfly and topological minors. It is natural to ask if we can drop this condition. This would require a different proof as ours draws upon this condition.

## References

- [1] Noga Alon. Piercing  $d$ -intervals. *Discrete & Computational Geometry*, 19(3):333–334, 1998.
- [2] Saeed Akhoondian Amiri, Ken-Ichi Kawarabayashi, Stephan Kreutzer, and Paul Wollan. The Erdős–Pósa property for directed graphs. *ArXiv e-prints*, 2016.
- [3] Julio Aracena, Jacques Demongeot, and Eric Goles. Positive and negative circuits in discrete neural networks. *Neural Networks, IEEE Transactions on*, 15(1):77–83, Jan 2004.
- [4] Julio Aracena, Adrien Richard, and Lilian Salinas. Number of fixed points and disjoint cycles in monotone Boolean networks. *ArXiv e-prints*, February 2016.
- [5] Maria Chudnovsky, Alexandra Fradkin, and Paul Seymour. Tournament immersion and cutwidth. *Journal of Combinatorial Theory, Series B*, 102(1):93 – 101, 2012.
- [6] Italo J. Dejter and Víctor Neumann-Lara. Unboundedness for generalized odd cyclic transversality. In *Combinatorics (Eger, 1987), Colloquia Mathematica Societatis János Bolyai*, volume 52, pages 195–203, 1987.
- [7] Paul Erdős and Lajos Pósa. On the maximal number of disjoint circuits of a graph. *Publicationes Mathematicae Debrecen*, 9:3–12, 1962.
- [8] Paul Erdős and Lajos Pósa. On independent circuits contained in a graph. *Canadian Journal of Mathematics*, 17:347–352, 1965.
- [9] Samuel Fiorini, Gwenaél Joret, and David R. Wood. Excluded forest minors and the Erdős–Pósa property. *Combinatorics, Probability & Computing*, 22(5):700–721, 2013.
- [10] Alexandra Fradkin and Paul Seymour. Tournament pathwidth and topological containment. *Journal of Combinatorial Theory, Series B*, 103(3):374 – 384, 2013.
- [11] Bertrand Guenin and Robin Thomas. Packing directed circuits exactly. *ArXiv e-prints*, December 2010.
- [12] András Gyárfás and Jenő Lehel. A Helly-type problem in trees. In *Combinatorial Theory and its applications*, pages 571–584. Colloquia Mathematica Societatis János Bolyai, 1969.



- [13] Frédéric Havet and Ana Karolinna Maia. On disjoint directed cycles with prescribed minimum lengths. Research Report RR-8286, INRIA, April 2013.
- [14] Thor Johnson, Neil Robertson, P.D. Seymour, and Robin Thomas. Directed tree-width. *Journal of Combinatorial Theory, Series B*, 82(1):138 – 154, 2001.
- [15] Ilhee Kim. *On containment relations in directed graphs*. PhD thesis, Princeton University, 2013.
- [16] Ilhee Kim and Paul Seymour. Tournament minors. *Journal of Combinatorial Theory, Series B*, 112:138 – 153, 2015.
- [17] Kenta Kitsunai, Yasuaki Kobayashi, and Hisao Tamaki. On the pathwidth of almost semicomplete digraphs. In Nikhil Bansal and Irene Finocchi, editors, *Algorithms - ESA 2015: 23rd Annual European Symposium, Patras, Greece, September 14-16, 2015, Proceedings*, pages 816–827, Berlin, Heidelberg, 2015. Springer Berlin Heidelberg.
- [18] Jean-Florent Raymond and Dimitrios M. Thilikos. Recent results on the Erdős–Pósa property. *Discrete Applied Mathematics*, to appear, 2017.
- [19] Bruce Reed. Mangoes and blueberries. *Combinatorica*, 19(2):267–296, 1999.
- [20] Bruce A. Reed. *Tree width and tangles: A new connectivity measure and some applications*, pages 87–162. Cambridge University Press, 1997.
- [21] Bruce A. Reed, Neil Robertson, Paul D. Seymour, and Robin Thomas. Packing directed circuits. *Combinatorica*, 16(4):535–554, 1996.
- [22] Bruce A. Reed and Bruce F. Shepherd. The Gallai-Younger conjecture for planar graphs. *Combinatorica*, 16(4):555–566, 1996.
- [23] Neil Robertson and Paul D. Seymour. Graph Minors. V. Excluding a planar graph. *Journal of Combinatorial Theory, Series B*, 41(2):92–114, 1986.
- [24] Paul D. Seymour. Packing circuits in Eulerian digraphs. *Combinatorica*, 16(2):223–231, 1996.