

Proofs Without Words

Exercises in Visual Thinking

© 1993 by
The Mathematical Association of America (Incorporated)
Library of Congress Catalog Card Number 93-86388

ISBN 0-88385-700-6

Printed in the United States of America

Current Printing (last digit):

10 9 8 7 6

Introduction

see (sē) *v.*, *saw*, *seen*, *seeing*. —*v.t.*

- ...
5. to perceive (things) mentally; discern;
understand: *to see the point of an argument*.

...
—THE RANDOM HOUSE DICTIONARY
OF THE ENGLISH LANGUAGE (2nd ED.)
UNABRIDGED.

"Proofs without words" (PWWs) have become regular features in the journals published by the Mathematical Association of America — notably *Mathematics Magazine* and *The College Mathematics Journal*. PWWs began to appear in *Mathematics Magazine* about 1975, and, in an editors' note in the January 1976 issue of the *Magazine*, J. Arthur Seebach and Lynn Arthur Steen encouraged further contributions of PWWs to the *Magazine*. Although originally solicited for "use as end-of-article fillers," the editors went on to ask "What could be better for this purpose than a pleasing illustration that made an important mathematical point?"

A few years earlier Martin Gardner, in his popular "Mathematical Games" column in the October 1973 issue of the *Scientific American*, discussed PWWs as "look-see" diagrams. Gardner points out that "in many cases a dull proof can be supplemented by a geometric analogue so simple and beautiful that the truth of a theorem is almost seen at a glance." This dramatically illustrates the dictionary quote above: in English "to see" is often "to understand."

In the same vein, the editorial policy of *The College Mathematics Journal* throughout most of the 1980s stated that, in addition to expository articles, "The Journal also invites other types of contributions, most notably: *proofs without words*, mathematical poetry, quotes, ..." (their italics). But PWWs are not recent innovations — they have a long history. Indeed, in this volume you will find modern renditions of proofs without words from ancient China, classical Greece, and India of the twelfth century.

Of course, “proofs without words” are not really proofs. As Theodore Eisenberg and Tommy Dreyfus note in their paper “On the Reluctance to Visualize in Mathematics” [in *Visualization in Teaching and Learning Mathematics*, MAA Notes Number 19], some consider such visual arguments to be of little value, and “that there is one and only one way to communicate mathematics, and ‘proofs without words’ are not acceptable.” But to counter this viewpoint, Eisenberg and Dreyfus go on to give us some quotes on the subject:

[Paul] Halmos, speaking of Solomon Lefschetz (editor of the *Annals*), stated: “He saw mathematics not as logic but as pictures.” Speaking of what it takes to be a mathematician, he stated: “To be a scholar of mathematics you must be born with ... the ability to visualize” and most teachers try to develop this ability in their students. [George] Pólya’s “Draw a figure ...” is classic pedagogic advice, and Einstein and Poincaré’s views that we should use our visual intuitions are well known.

So, if “proofs without words” are not proofs, what are they? As you will see from this collection, this question does not have a simple, concise answer. But generally, PWWs are pictures or diagrams that help the observer see *why* a particular statement may be true, and also to see *how* one might begin to go about proving it true. In some an equation or two may appear in order to guide the observer in this process. But the emphasis is clearly on providing visual clues to the observer to stimulate mathematical thought.

I should note that this collection is not intended to be complete. It does not include all PWWs which have appeared in print, but is rather a sample representative of the genre. In addition, as readers of the Association’s journals are well aware, new PWWs appear in print rather frequently, and I anticipate that this will continue. Perhaps some day a second volume of PWWs will appear!

I hope that the readers of this collection will find enjoyment in discovering or rediscovering some elegant visual demonstrations of certain mathematical ideas; that teachers will want to share many of them with their students; and that all will find stimulation and encouragement to try to create new “proofs without words.”

Acknowledgment. I would like to express my appreciation and gratitude to the many people who have played a part in the publication of this collection: to Gerald Alexanderson and Martha Siegel, who, as editors of *Mathematics Magazine*, gave me encouragement over the years as I learned to read and write PWWs; to Doris Schattschneider, Eugene Klotz, and Richard Guy for sharing with me their collections of PWWs; and finally, to all those individuals who have contributed “proofs without words” to the mathematical literature (see the *Index of Names* on pp. 151-152), without whom this collection simply would not exist.

Note. All the drawings in this collection were redone to create a uniform appearance. In a few instances titles were changed, and shading or symbols were added (or deleted) for clarity. Any errors resulting from that process are entirely my responsibility.

Roger B. Nelsen
Lewis and Clark College
Portland, Oregon



Contents

Introduction	v
Geometry & Algebra	1
Trigonometry, Calculus & Analytic Geometry	27
Inequalities	47
Integer Sums	67
Sequences & Series	113
Miscellaneous	131
Sources	145
Index of Names	151

Geometry & Algebra



The Pythagorean Theorem I	3
The Pythagorean Theorem II	4
The Pythagorean Theorem III	5
The Pythagorean Theorem IV	6
The Pythagorean Theorem V	7
The Pythagorean Theorem VI	8
A Pythagorean Theorem: $a \cdot a' = b \cdot b' + c \cdot c'$	9
The Rolling Circle Squares Itself	10
On Trisecting an Angle	11
Trisection of an Angle in an Infinite Number of Steps	12
Trisection of a Line Segment	13
The Vertex Angles of a Star Sum to 180°	14
Viviani's Theorem	15
A Theorem About Right Triangles	16
Area and the Projection Theorem of a Right Triangle	17
Chords and Tangents of Equal Length	18
Completing the Square	19
Algebraic Areas I	20
Algebraic Areas II	21
Diophantus of Alexandria's "Sum of Squares" Identity	22
The k^{th} n -gonal Number	23
The Volume of a Frustum of a Square Pyramid	24
The Volume of a Hemisphere via Cavalieri's Principle	25

The Pythagorean Theorem I



—adapted from the *Chou pei suan ching*
(author unknown, circa B.C. 200?)



The Pythagorean Theorem II



Behold!


—Bhāskara (12th century)

The Pythagorean Theorem III




—based on Euclid's proof

The Pythagorean Theorem IV



—H. E. Dudeney (1917)


The Pythagorean Theorem V



—James A. Garfield (1876)
20th President of the United States



The Pythagorean Theorem VI

$$\frac{c+a}{b} = \frac{b}{c-a}$$
$$a^2 + b^2 = c^2$$



—Michael Hardy

A Pythagorean Theorem: $a \cdot a' = b \cdot b' + c \cdot c'$




$$\frac{x}{b} = \frac{b'}{a} \Rightarrow a \cdot x = b \cdot b';$$

$$\frac{y}{c} = \frac{c'}{a} \Rightarrow a \cdot y = c \cdot c';$$


$$\therefore a \cdot a' = a \cdot (x + y) = b \cdot b' + c \cdot c'.$$

The Rolling Circle Squares Itself




—Thomas Elsner

On Trisecting an Angle



—Rufus Isaacs





Trisection of an Angle in an Infinite Number of Steps



$$\frac{1}{3} = \frac{1}{2} - \frac{1}{4} + \frac{1}{8} - \frac{1}{16} + \dots$$


—Eric Kincanon

Trisection of a Line Segment




$$\overline{AF} = \frac{1}{3} \cdot \overline{AB}$$

The Vertex Angles of a Star Sum to 180°



Viviani's Theorem


The perpendiculars to the sides from a point on the boundary or within an equilateral triangle add up to the height of the triangle.



—Samuel Wolf


A Theorem About Right Triangles

The internal bisector of the right angle of a right triangle bisects the square on the hypotenuse.



—Roland H. Eddy

Area and the Projection Theorem of a Right Triangle




$$CD^2 = AD \cdot DB$$

—Sidney H. Kung


Chords and Tangents of Equal Length

If circle C_1 passes through the center O of circle C_2 , the length of the common chord \overline{PQ} is equal to the tangent segment \overline{PR} .




Completing the Square

$$x^2 + ax = (x + a/2)^2 - (a/2)^2$$





Algebraic Areas I

$$(a + b)^2 + (a - b)^2 = 2(a^2 + b^2)$$




Algebraic Areas II

$$(a + b + c)^2 + (a + b - c)^2 + (a - b + c)^2 + (a - b - c)^2 \\ = (2a)^2 + (2b)^2 + (2c)^2$$




Diophantus of Alexandria's "Sum of Squares" Identity

$$(a^2 + b^2)(c^2 + d^2) = (ad + bc)^2 + (bd - ac)^2$$






The k th n -gonal Number is

$$1 + (k - 1)(n - 1) + \frac{1}{2}(k - 2)(k - 1)(n - 2)$$



The Volume of a Frustum of a Square Pyramid

[Problem 14, *The Moscow Papyrus*, circa 1850 b.c.]


 P_1  P_2  $3P_2$

$$V(P_1) = \frac{h}{b-a} V(P_2) = \frac{h}{b-a} \cdot \frac{1}{3} (b^3 - a^3) = \frac{h}{3} (a^2 + ab + b^2)$$

REFERENCES

1. C. B. Boyer, *A History of Mathematics*, John Wiley & Sons, New York, 1968, pp. 20-22.
2. R. J. Gillings, *Mathematics in the Time of the Pharaohs*, The MIT Press, Cambridge, 1972, pp. 187-193.

The Volume of a Hemisphere via Cavalieri's Principle*



$$V_S = V_P = \frac{1}{3} r^2 \cdot 2\pi r = \frac{2}{3} \pi r^3$$

*Tzu Geng, son of the most celebrated mathematician Tzu Chung Chih in ancient China, was believed to be the first to develop the principle in the 5th century A. D.


—Sidney H. Kung

Trigonometry, Calculus & Analytic Geometry

Sine of the Sum	29	
Area and Difference Formulas	30	
The Law of Cosines I	31	
The Law of Cosines II	32	
The Law of Cosines III (via Ptolemy's Theorem)	33	
The Double-Angle Formulas	34	
The Half-Angle Tangent Formulas	35	
Mollweide's Equation	36	
$(\tan\theta + 1)^2 + (\cot\theta + 1)^2 = (\sec\theta + \csc\theta)^2$	37	
The Substitution to Make a Rational Function of the Sine and Cosine		38
Sums of Arctangents	39	
The Distance Between a Point and a Line	40	
The Midpoint Rule is Better than the Trapezoidal Rule for Concave Functions		41
Integration by Parts	42	
The Graphs of f and f^{-1} are Reflections about the Line $y = x$	43	
The Reflection Property of the Parabola	44	
Area under an Arch of the Cycloid	45	

Sine of the Sum

$$\sin(x+y) = \sin x \cos y + \cos x \sin y \text{ for } x+y < \pi$$




$$c = a \cos y + b \cos x$$


$$r = 1/2 \Rightarrow \sin z = (c/2)/(1/2) = c, \sin x = a, \sin y = b;$$

$$\sin(x+y) = \sin(\pi - (x+y)) = \sin z = \sin x \cos y + \sin y \cos x$$

Area and Difference Formulas




$$\sin(x - y) = \sin x \cos y - \cos x \sin y$$



$$\cos(x - y) = \cos x \cos y + \sin x \sin y$$


—Sidney H. Kung

The Law of Cosines I



$$\begin{aligned}c^2 &= (b \sin \theta)^2 + (a - b \cos \theta)^2 \\&= a^2 + b^2 - 2ab \cos \theta\end{aligned}$$

The Law of Cosines II




$$(2a \cos \theta - b)b = (a - c)(a + c)$$

$$c^2 = a^2 + b^2 - 2ab \cos \theta$$

—Sidney H. Kung


The Law Of Cosines III (via Ptolemy's Theorem)



$$c \cdot c = b \cdot b + (a + 2b \cos(\pi - \theta)) \cdot a$$

$$c^2 = a^2 + b^2 - 2ab \cdot \cos\theta$$

The Double-Angle Formulas



$$\Delta ACD \sim \Delta ABC$$

$$\frac{CD}{AC} = \frac{BC}{AB}$$

$$\frac{AD}{AC} = \frac{AC}{AB}$$


$$\frac{\sin 2\theta}{2\cos\theta} = \frac{2\sin\theta}{2}$$

$$\frac{(1 + \cos 2\theta)}{2\cos\theta} = \frac{2\cos\theta}{2}$$

$$\sin 2\theta = 2\sin\theta\cos\theta$$

$$\cos 2\theta = 2\cos^2\theta - 1$$


The Half-Angle Tangent Formulas



$$\tan \frac{\theta}{2} = \frac{\sin\theta}{1 + \cos\theta} = \frac{1 - \cos\theta}{\sin\theta}$$


Mollweide's Equation

$$(a - b) \cos \frac{\gamma}{2} = c \sin \left(\frac{\alpha - \beta}{2} \right)$$



—H. Arthur DeKleine

$$(\tan\theta + 1)^2 + (\cot\theta + 1)^2 = (\sec\theta + \csc\theta)^2$$




$$\tan^2\theta + 1 = \sec^2\theta$$

$$\cot^2\theta + 1 = \csc^2\theta$$

$$(\tan\theta + 1)^2 + (\cot\theta + 1)^2 = (\sec\theta + \csc\theta)^2$$


$$\left(\text{also } \tan\theta = \frac{\tan\theta + 1}{\cot\theta + 1} \right)$$

The Substitution to Make a Rational Function of the Sine and Cosine




$$z = \tan \frac{\theta}{2} \Rightarrow \sin \theta = \frac{2z}{1+z^2} \text{ and } \cos \theta = \frac{1-z^2}{1+z^2}$$

Sums of Arctangents




$$\arctan \frac{1}{2} + \arctan \frac{1}{3} = \frac{\pi}{4}$$



$$\arctan 1 + \arctan 2 + \arctan 3 = \pi$$

—Edward M. Harris


The Distance Between a Point and a Line




$$\frac{d}{1} = \frac{|ma + c - b|}{\sqrt{1 + m^2}}$$

—R. L. Eisenman

The Midpoint Rule is Better than the Trapezoidal Rule for Concave Functions



Integration by Parts





$$\text{Area } \boxed{\text{white}} + \text{Area } \boxed{\text{hatched}} = qs - pr$$

$$\int\limits_r^s u \, dv + \int\limits_p^q v \, du = uv \Big|_{(p,r)}^{(q,s)}$$


$$\int\limits_a^b f(x)g'(x)dx = f(x)g(x) \Big|_a^b - \int\limits_a^b g(x)f'(x)dx$$

—Richard Courant

The Graphs of f and f^{-1} are Reflections about the Line $y = x$






The Reflection Property of the Parabola



$$QF = QD \quad \& \quad m_1 \cdot m_2 = -1 \quad \Rightarrow \quad \angle 1 = \angle 2 = \angle 3$$

—Ayoub B. Ayoub

Area under an Arch of the Cycloid




$$\frac{1}{2} \pi R \cdot 2R + \pi R^2 + \frac{1}{2} \pi R \cdot 2R \\ \Rightarrow A = 3 \pi R^2$$

—Richard M. Beekman

Inequalities


The Arithmetic Mean—Geometric Mean Inequality I	49
The Arithmetic Mean—Geometric Mean Inequality II	50
The Arithmetic Mean—Geometric Mean Inequality III	51
Two Extremum Problems	52
The Harmonic Mean—Geometric Mean—Arithmetic Mean— Root Mean Square Inequality I	53
The Harmonic Mean—Geometric Mean—Arithmetic Mean— Root Mean Square Inequality II	54
The Harmonic Mean—Geometric Mean—Arithmetic Mean— Root Mean Square Inequality III	55
Five Means — and Their Means	56
$e^\pi > \pi^e$	58
$A^B > B^A$ for $e \leq A < B$	59
The Mediant Property	60
Regle des Nombres Moyens	61
The Sum of a Positive Number and its Reciprocal is at least Two	62
Aristarchus' Inequalities	63
The Cauchy-Schwarz Inequality	64
Bernoulli's Inequality	65
Napier's Inequality	66

The Arithmetic Mean—Geometric Mean Inequality I



$$\sqrt{ab} \leq \frac{a+b}{2}$$

The Arithmetic Mean—Geometric Mean Inequality II




$$(a+b)^2 - (a-b)^2 = 4ab$$

$$\frac{a+b}{2} \geq \sqrt{ab}$$

—Doris Schattschneider


The Arithmetic Mean—Geometric Mean Inequality III

$$\frac{a+b}{2} \geq \sqrt{ab}, \text{ with equality if and only if } a = b$$




Two Extremum Problems


For a given product, the sum of two positive numbers is minimal when the numbers are equal.



For a given sum, the product of two positive numbers is maximal when the numbers are equal.



The Harmonic Mean—Geometric Mean—
 Arithmetic Mean—Root Mean Square
 Inequality I




$$PM = a, QM = b, a > b > 0$$

$$HM < GM < AM < RM$$

$$\frac{2ab}{a+b} < \sqrt{ab} < \frac{a+b}{2} < \sqrt{(a^2 + b^2)/2}$$


The Harmonic Mean—Geometric Mean—
 Arithmetic Mean—Root Mean Square
 Inequality II



—Sidney H. Kung


The Harmonic Mean—Geometric Mean—Arithmetic Mean—Root Mean Square Inequality III

$$a, b > 0 \Rightarrow \sqrt{(a^2 + b^2)/2} \geq \frac{a+b}{2} \geq \sqrt{ab} \geq \frac{2ab}{a+b}$$




$$2a^2 + 2b^2 \geq (a+b)^2$$

$$\sqrt{\frac{a^2 + b^2}{2}} \geq \frac{a+b}{2}$$



$$(\sqrt{a+b})^2 \geq 4 \cdot \frac{1}{2} \sqrt{a} \sqrt{b}$$

$$\frac{a+b}{2} \geq \sqrt{ab}$$



$$1 \geq 4 \frac{a}{a+b} \cdot \frac{b}{a+b}$$

$$\sqrt{ab} \geq \frac{2ab}{a+b}$$

Five Means — and Their Means


Arithmetic: $am = AM(a,b) = \frac{a+b}{2}$

Contraharmonic: $cm = CM(a,b) = \frac{a^2 + b^2}{a+b}$

Geometric: $gm = GM(a,b) = \sqrt{ab}$

Harmonic: $hm = HM(a,b) = \frac{2ab}{a+b}$

Root Mean Square: $rms = RMS(a,b) = \sqrt{\frac{a^2 + b^2}{2}}$



I. $0 < a < b \Rightarrow$

$$a < \frac{2ab}{a+b} < \sqrt{ab} < \frac{a+b}{2} < \sqrt{\frac{a^2+b^2}{2}} < \frac{a^2+b^2}{a+b} < b$$

II. $hm + cm = a + b \Rightarrow AM(hm, cm) = am.$


III. $hm \cdot am = a \cdot b \Rightarrow GM(hm, am) = gm.$

IV. $am \cdot cm = \frac{a^2 + b^2}{2} \Rightarrow GM(am, cm) = rms.$

V. $gm^2 + rms^2 = \frac{(a+b)^2}{2} \Rightarrow RMS(gm, rms) = am.$


$$a < \frac{2ab}{a+b} < \sqrt{ab} < \frac{a+b}{2} < \sqrt{\frac{a^2+b^2}{2}} < \frac{a^2+b^2}{a+b} < b$$

$$e^\pi > \pi^e$$



—Fouad Nakhli

$A^B > B^A$ for $e \leq A < B$




$$e \leq A < B \Rightarrow m_A > m_B$$

$$\Rightarrow \frac{\ln A}{A} > \frac{\ln B}{B}$$

$$\Rightarrow A^B > B^A$$

The Mediant Property

$$\frac{a}{b} < \frac{c}{d} \Rightarrow \frac{a}{b} < \frac{a+c}{b+d} < \frac{c}{d}$$




Regle des Nombres Moyens (Two Proofs)

[Nicolas Chuquet, *Le Triparty en la Science des Nombres*, 1484]

$$a, b, c, d > 0; \frac{a}{b} < \frac{c}{d} \Rightarrow \frac{a}{b} < \frac{a+c}{b+d} < \frac{c}{d}$$


I.



$$m_1 < m_3 \Rightarrow m_1 < m_2 < m_3$$

—Li Changming

II.




$$\frac{a}{b} < \frac{a}{b+d} + \frac{c}{b+d} < \frac{c}{d}$$


—RBN

The Sum of a Positive Number and its Reciprocal is at least Two (Four Proofs)


I.




II.



III.




IV.



$$x \geq 1 \Rightarrow x + \frac{1}{x} \geq 2$$

Aristarchus' Inequalities

$$0 < \beta < \alpha < \frac{\pi}{2} \Rightarrow \frac{\sin \alpha}{\sin \beta} < \frac{\alpha}{\beta} < \frac{\tan \alpha}{\tan \beta}$$




$$\sin \alpha < \frac{\sin \beta}{\beta} \alpha; \quad \frac{\tan \beta}{\beta} \alpha < \tan \alpha$$

$$\therefore \frac{\sin \alpha}{\sin \beta} < \frac{\alpha}{\beta} < \frac{\tan \alpha}{\tan \beta}$$

The Cauchy-Schwarz Inequality

$$|\langle a, b \rangle \cdot \langle x, y \rangle| \leq \|\langle a, b \rangle\| \|\langle x, y \rangle\|$$




$$(|a| + |b|)(|b| + |x|) \leq 2\left(\frac{1}{2}|a||b| + \frac{1}{2}|x||y|\right) + \sqrt{a^2 + b^2} \sqrt{x^2 + y^2}$$

$$\therefore |ax + by| \leq |a||x| + |b||y| \leq \sqrt{a^2 + b^2} \sqrt{x^2 + y^2}$$


Bernoulli's Inequality (two proofs)

$$x > 0, x \neq 1, r > 1 \Rightarrow x^r - 1 > r(x - 1)$$


I. (first semester calculus)



II. (second semester calculus)



$$x^r - 1 = \int_1^x rt^{r-1} dt > r(x - 1)$$




$$1 - x^r = \int_x^1 rt^{r-1} dt < r(1 - x)$$


Napier's Inequality (two proofs)

$$b > a > 0 \Rightarrow \frac{1}{b} < \frac{\ln b - \ln a}{b - a} < \frac{1}{a}$$

I. (first semester calculus)



II. (second semester calculus)




$$\frac{1}{b}(b-a) < \int_a^b \frac{1}{x} dx < \frac{1}{a}(b-a)$$

Integer Sums

Sums of Integers I	69
Sums of Integers II	70
Sums of Odd Integers I	71
Sums of Odd Integers II	72
Sums of Odd Integers III	73
Squares and Sums of Integers	74
Arithmetic Progressions with Sum Equal to the Square of the Number of Terms	76
Sums of Squares I	77
Sums of Squares II	78
Sums of Squares III	79
Sums of Squares IV	80
Sums of Squares V	81
Alternating Sums of Squares	82
Sums of Squares of Fibonacci Numbers	83
Sums of Cubes I	84
Sums of Cubes II	85
Sums of Cubes III	86
Sums of Cubes IV	87
Sums of Cubes V	88
Sums of Cubes VI	89
Sums of Integers and Sums of Cubes	90
Sums of Odd Cubes are Triangular Numbers	91

Sums of Fourth Powers	92
k^{th} Powers as Sums of Consecutive Odd Numbers	93
Sums of Triangular Numbers I	94
Sums of Triangular Numbers II	95
Sums of Triangular Numbers III	96
Sums of Oblong Numbers I	97
Sums of Oblong Numbers II	98
Sums of Oblong Numbers III	99
Sums of Pentagonal Numbers	100
On Squares of Positive Integers	101
Consecutive Sums of Consecutive Integers	102
Count the Dots	103
Identities for Triangular Numbers	104
A Triangular Identity	105
Every Hexagonal Number is a Triangular Number	106
One Domino = Two Squares: Concentric Squares	107
Sums of Consecutive Powers of Nine are	
Sums of Consecutive Integers	108
Sums of Hex Numbers Are Cubes	109
Every Cube is the Sum of Consecutive Odd Numbers	110
The Cube as an Arithmetic Sum	111


Sums of Integers I



$$1 + 2 + \dots + n = \frac{1}{2}n(n + 1)$$

—“The ancient Greeks”
(as cited by Martin Gardner)


Sums of Integers II



$$1 + 2 + \cdots + n = \frac{n^2}{2} + \frac{n}{2}$$

—Ian Richards


Sums of Odd Integers I



$$1 + 3 + 5 + \cdots + (2n - 1) = n^2$$


—Nicomachus of Gerasa (circa A.D. 100)

Sums of Odd Integers II



$$1 + 3 + \cdots + (2n - 1) = \frac{1}{4}(2n)^2 = n^2$$

Sums of Odd Integers III




$$\Delta + 3\cdot\Delta + \dots + (2n-1)\cdot\Delta = A = n^2\cdot\Delta$$


$$\sum_{i=1}^n (2i-1) = n^2$$

Squares and Sums of Integers


I.



$$1 + 2 + 1 = 2^2$$



$$1 + 2 + 3 + 2 + 1 = 3^2$$




$$1 + 2 + 3 + 4 + 3 + 2 + 1 = 4^2$$


$$1 + 2 + \dots + (n-1) + n + (n-1) + \dots + 2 + 1 = n^2$$

—“The ancient Greeks”
(as cited by Martin Gardner)

II.




$$= \bullet +$$




$$1 + 3 + 1 = 1^2 + 2^2$$




$$= \bullet \bullet +$$



$$1 + 3 + 5 + 3 + 1 = 2^2 + 3^2$$



$$= \bullet \bullet \bullet +$$



$$1 + 3 + 5 + 7 + 5 + 3 + 1 = 3^2 + 4^2$$


•

•




•

$$1 + 3 + \dots + (2n-1) + (2n+1) + (2n-1) + \dots + 3 + 1 = n^2 + (n+1)^2$$

Arithmetic Progressions with Sum Equal to the Square of the Number of Terms



$$\sum_{k=n}^{3n-2} k = (2n-1)^2; \quad n=1, 2, 3, \dots.$$





$$n = 4$$

$$4 + 5 + 6 + 7 + 8 + 9 + 10 = 7^2$$

—James O. Chilaka




Sums of Squares I

$$1^2 + 2^2 + \dots + n^2 = \frac{1}{3}n(n+1)(n+\frac{1}{2})$$




Sums of Squares II

$$3(1^2 + 2^2 + \dots + n^2) = (2n + 1)(1 + 2 + \dots + n)$$



1 + 2 + ⋯ + n



$2n + 1$

—Martin Gardner and Dan Kalman
(independently)

Sums of Squares III


$$3(1^2 + 2^2 + \dots + n^2) = \frac{1}{2}n(n+1)(2n+1)$$

$$\begin{array}{ccccccccc}
 n & n & \cdots & n & n & n & n-1 & \cdots & 2 & 1 & 1 & 2 & \cdots & n-1 & n \\
 n-1 & n-1 & \cdots & n-1 & n & n-1 & \cdots & 2 & & 2 & 3 & \cdots & n \\
 \cdot & & \cdot & \cdot & \cdot & \cdot \\
 \cdot & \cdot & \cdot & & + & \cdot & \cdot & \cdot & & + & \cdot & \cdot & \cdot \\
 \cdot & \cdot & \cdot & & & \cdot & \cdot & \cdot & & & \cdot & \cdot & \cdot \\
 2 & 2 & & & n & n-1 & & & & n-1 & n & & \\
 1 & & & & n & & & & & n & & &
 \end{array}$$

$$\begin{array}{cccccc}
 & & & & & \\
 & 2n+1 & 2n+1 & \cdots & 2n+1 & 2n+1 \\
 & 2n+1 & 2n+1 & \cdots & 2n+1 & \\
 & \cdot & \cdot & & \cdot & \\
 = & \cdot & \cdot & \cdot & \cdot & \\
 & \cdot & \cdot & \cdot & & \\
 & 2n+1 & 2n+1 & & & \\
 & 2n+1 & & & &
 \end{array}$$

Sums of Squares IV


$$\sum_{k=1}^n k^2 = \left(\sum_{k=1}^n k \right)^2 - 2 \sum_{k=1}^{n-1} \left[\left(\sum_{i=1}^k i \right) (k+1) \right]$$



—James O. Chilaka


Sums of Squares V

$$\sum_{i=1}^n \sum_{j=i}^n j = \sum_{i=1}^n i^2$$



Alternating Sums of Squares


I.



$$\sum_{k=1}^n (-1)^{k+1} k^2 = (-1)^{n+1} T_n = (-1)^{n+1} \frac{n(n+1)}{2}$$

—Dave Logothetti


II.



$$n^2 - (n-1)^2 + \dots + (-1)^{n-1} (1)^2 = \sum_{k=0}^n (-1)^k (n-k)^2 = \frac{n(n+1)}{2}$$

—Steven L. Snover

Sums of Squares of Fibonacci Numbers




$$F_1 = F_2 = 1; F_{n+2} = F_{n+1} + F_n \Rightarrow F_1^2 + F_2^2 + \dots + F_n^2 = F_n F_{n+1}$$

—Alfred Brousseau

Sums of Cubes I


$$1^3 + 2^3 + 3^3 + \cdots + n^3 = (1 + 2 + 3 + \cdots + n)^2$$



—Solomon W. Golomb

Sums of Cubes II










$$1^3 + 2^3 + 3^3 + \cdots + n^3 = (1 + 2 + 3 + \cdots + n)^2$$



—J. Barry Love

Sums of Cubes III


$$1^3 + 2^3 + 3^3 + \cdots + n^3 = (1 + 2 + 3 + \cdots + n)^2$$



—Alan L. Fry


Sums of Cubes IV

$$1^3 + 2^3 + 3^3 + \cdots + n^3 = \frac{1}{4}[n(n+1)]^2$$




—Antonella Cupillari and Warren Lushbaugh
(independently)

Sum of Cubes V



$$t_n = 1 + 2 + \cdots + n \Rightarrow t_n^2 - t_{n-1}^2 = n^3$$



$$t_n^2 = (1 + 2 + \cdots + n)^2 = 1^3 + 2^3 + 3^3 + \cdots + n^3$$

Sum of Cubes VI

	1	2	3	.	.	.	n
+	2	4	6	.	.	.	2n
+	3	6	9	.	.	.	3n
+
.
.
+	n	2n	3n	.	.	.	n ²

	1	2	3	.	.	.	n
+	2	4	6	.	.	.	2n
+	3	6	9	.	.	.	3n
+
.
.
+	n	2n	3n	.	.	.	n ²

$$= \sum_{i=1}^n i + 2 \sum_{i=1}^n i + \dots + n \sum_{i=1}^n i$$

$$= 1(1)^2 + 2(2)^2 + \dots + n(n)^2$$


$$= \left(\sum_{i=1}^n i \right)^2$$

$$= \sum_{i=1}^n i^3$$

Sums of Integers and Sums of Cubes

$$1 + 2 + \cdots + n = \frac{1}{2}n(n + 1)$$

$$1^3 + 2^3 + \cdots + n^3 = \left(\frac{1}{2}n(n + 1) \right)^2$$



—Georg Schrage

Sums of Odd Cubes are Triangular Numbers

$$1^3 = \square$$

$$3^3 = 3(3^2) = \boxed{9} + \boxed{9} + \boxed{9} = \begin{array}{c} \text{A 3x3 grid of squares. The first two columns are shaded black, and the third column is white. A double-headed arrow below the grid indicates its width is } 1+2(3). \\ \hline \end{array}$$

$$5^3 = 5(5^2) = \boxed{\text{grid}} + \boxed{\text{grid}} + \boxed{\text{grid}}$$

$$\leftarrow 2 + 3(5) \rightarrow$$

$$(2n-1)^3 = (2n-1)(2n-1)^2 = \dots$$


A diagram illustrating a geometric proof. A large right-angled triangle is divided into n^2 smaller triangles by a zigzag line from the bottom-left vertex to the top-right vertex. The base of the triangle is labeled with the expression $(n - 1) + n(2n - 1)$.

$$1^3 + 3^3 + 5^3 + \dots + (2n-1)^3 = 1 + 2 + 3 + \dots + (2n^2-1) = n^2(2n^2-1)$$

—Monte J. Zerger

Sums of Fourth Powers


$$\sum_{i=1}^n i^4 = \left(\sum_{i=1}^n i^2 \right)^2 - 2 \left[\sum_{k=2}^n \left(k^2 \sum_{i=1}^{k-1} i^2 \right) \right]$$



—Elizabeth M. Markham


k^{th} Powers as Sums of Consecutive Odd Numbers

$$n^k = (n^{k-1} - n + 1) + (n^{k-1} - n + 3) + \dots + (n^{k-1} - n + 2n - 1); \\ k = 2, 3, \dots.$$



Sums of Triangular Numbers I

$$T_n = 1 + 2 + \cdots + n \Rightarrow T_1 + T_2 + \cdots + T_n = \frac{n(n+1)(n+2)}{6}$$




$$3(T_1 + T_2 + \cdots + T_n) = (n+2) \cdot T_n$$

$$T_1 + T_2 + \cdots + T_n = \frac{(n+2)}{3} \cdot \frac{n(n+1)}{2} = \frac{n(n+1)(n+2)}{6}$$

Sums of Triangular Numbers II

$$T_k = 1 + 2 + \cdots + k \Rightarrow \sum_{k=1}^n T_k = \frac{1}{6} n(n+1)(n+2)$$



Sums of Triangular Numbers III

$$T_k = 1 + 2 + \cdots + k \Rightarrow 3 \sum_{k=1}^n T_k = \frac{1}{2} n(n+1)(n+2)$$

$$\begin{array}{ccccccc}
 1 & & 1 & & & & n \\
 1 2 & & 2 1 & & & & n-1 n-1 \\
 1 2 3 & & 3 2 1 & & & & n-2 n-2 n-2 \\
 \cdot \cdot \cdot & + & \cdot \cdot \cdot & + & \cdot \cdot \cdot & & \cdot \cdot \cdot \\
 \cdot \cdot \cdot & & \cdot \cdot \cdot & & & & \cdot \cdot \cdot \\
 1 2 \cdots n-1 & & n-1 n-2 & \cdots & 1 & 2 2 & \cdots 2 \\
 1 2 \cdots n-1 n & & n n-1 & \cdots & 2 1 & 1 1 & \cdots 1 1
 \end{array}$$


$$\begin{array}{c}
 n+2 \\
 n+2 n+2 \\
 n+2 n+2 n+2 \\
 = \cdot \cdot \cdot \\
 \cdot \cdot \cdot \\
 n+2 n+2 \cdots n+2 \\
 n+2 n+2 \cdots n+2 n+2
 \end{array}$$

$$3(T_1 + T_2 + \cdots + T_n) = T_n \cdot (n+2)$$


Sums of Oblong Numbers I

$$(1 \times 2) + (2 \times 3) + (3 \times 4) + \cdots + (n-1)n = \frac{(n-1)n(n+1)}{3}$$

(i)





(ii)



Sums of Oblong Numbers II



$$3(1 \cdot 2 + 2 \cdot 3 + 3 \cdot 4 + \cdots + n(n+1)) = n(n+1)(n+2)$$






—Sidney H. Kung

Sums of Oblong Numbers III




$$(1 \times 2) + (2 \times 3) + \cdots + (n - 1) \times n = \frac{1}{3}[n^3 - n]$$





$$3(1 \times 2) = 2^3 - 2$$



$$3(1 \times 2) + 3(2 \times 3) = 3^3 - 3$$




$$3(1 \times 2) + 3(2 \times 3) + 3(3 \times 4) =$$



$$4^3 - 4$$

Sums of Pentagonal Numbers


$$\frac{1 \cdot 2}{2} + \frac{2 \cdot 5}{2} + \frac{3 \cdot 8}{2} + \cdots + \frac{n(3n - 1)}{2} = \frac{n^2(n + 1)}{2}$$




—William A. Miller

On Squares of Positive Integers

$$T_n = 1 + 2 + \cdots + n \Rightarrow$$




$$(2n+1)^2 = 8T_n + 1$$



$$(2n)^2 = 8T_{n-1} + 4n$$


Consecutive Sums of Consecutive Integers




$$\begin{aligned}
 1 + 2 &= 3 \\
 4 + 5 + 6 &= 7 + 8 \\
 9 + 10 + 11 + 12 &= 13 + 14 + 15 \\
 16 + 17 + 18 + 19 + 20 &= 21 + 22 + 23 + 24
 \end{aligned}$$

$$n^2 + (n^2 + 1) + \dots + (n^2 + n) = (n^2 + n + 1) + \dots + (n^2 + 2n)$$

Count the Dots




$$\sum_{k=1}^n k + \sum_{k=1}^{n-1} k = n^2$$



$$\sum_{k=1}^n k + n^2 = \sum_{k=n+1}^{2n} k$$


Identities for Triangular Numbers

$$T_n = 1 + 2 + \cdots + n \Rightarrow$$




$$3T_n + T_{n-1} = T_{2n}$$

$$3T_n + T_{n+1} = T_{2n+1}$$



$$T_{n-1} + 6T_n + T_{n+1} = (2n+1)^2$$


A Triangular Identity




$$T_n = 1 + 2 + \dots + n \Rightarrow (2k+1)^2 T_n + T_k = T_{(2k+1)n+k}$$

Every Hexagonal Number is a Triangular Number


$$\left. \begin{array}{l} H_n = 1+5+\dots+(4n-3) \\ T_n = 1+2+\dots+n \end{array} \right\} \Rightarrow H_n = 3T_{n-1} + T_n = T_{2n-1} = n(2n-1)$$



H_5




T_9



5.9

One Domino = Two Squares: Concentric Squares









$$1 + 4 \cdot 2 + 8 \cdot 2 + 12 \cdot 2 + 16 \cdot 2 = 9^2$$

$$1 + 2 \sum_{k=1}^n 4k = (2n+1)^2$$

—Shirley A. Wakin


Sums of Consecutive Powers of Nine are Sums of Consecutive Integers



$$1 + 9 + \dots + 9^n = 1 + 2 + 3 + \dots + (1 + 3 + \dots + 3^n)$$

—RBN

Sums of Hex Numbers Are Cubes





$$h_1 = 1$$

$$h_2 = 7$$

$$h_3 = 19$$


$$h_4 = 37$$




$$h_n = n^3 - (n-1)^3$$

$$\therefore h_1 + h_2 + \cdots + h_n = n^3.$$







Every Cube is the Sum of Consecutive Odd Numbers



The Cube as an Arithmetic Sum



$$n^3 = \sum_{i=0}^{n-1} 2i(n+1) + 1$$



$$5^3 = 1 + 13 + 25 + 37 + 49$$


—Robert Bronson and
Christopher Brueningsen

Sequences & Series

On a Property of the Sequence of Odd Integers (Galileo, 1615)	115
A Monotone Sequence Bounded by e	116
A Recursively Defined Sequence for e	117
Geometric Sums	118
Geometric Series I	119
Geometric Series II	120
Geometric Series III	121
Geometric Series IV	122
Gabriel's Staircase	123
Differentiated Geometric Series	124
$\frac{1}{1\cdot2} + \frac{1}{2\cdot3} + \cdots + \frac{1}{n(n+1)} = \frac{n}{n+1}$	126
The Series of Reciprocals of Triangular Numbers	127
The Alternating Harmonic Series	128
$\sin(2n + 1)\theta = \sin\theta + 2\sin\theta \sum_{k=1}^n \cos 2k\theta$	129
An Arctangent Identity and Series	130

On a Property of the Sequence of Odd Integers (Galileo, 1615)

$$\frac{1}{3} = \frac{1+3}{5+7} = \frac{1+3+5}{7+9+11} = \dots .$$




$$\frac{1 + 3 + \dots + (2n - 1)}{(2n + 1) + (2n + 3) + \dots + (4n - 1)} = \frac{1}{3}$$

REFERENCE

S. Drake, *Galileo Studies*, The University of Michigan Press, Ann Arbor, 1970, pp. 218-219.

A Monotone Sequence Bounded by e

$$\forall n \geq 1, \left(1 + \frac{1}{n}\right)^n < \left(1 + \frac{1}{n+1}\right)^{n+1} < e.$$




$$n \geq 1 \Rightarrow m_1 < m_2 < 1$$

$$\Rightarrow \frac{\ln\left(1 + \frac{1}{n}\right)}{\frac{1}{n}} < \frac{\ln\left(1 + \frac{1}{n+1}\right)}{\frac{1}{n+1}} < 1$$

$$\Rightarrow \left(1 + \frac{1}{n}\right)^n < \left(1 + \frac{1}{n+1}\right)^{n+1} < e$$


A Recursively Defined Sequence for e




$$x_0 > 1 \text{ } \& \text{ } x_{n+1} = \frac{x_n}{\ln(x_n)} \Rightarrow \lim x_n = e$$

—Thomas P. Dence

Geometric Sums





$$\frac{1}{2} + \frac{1}{4} + \frac{1}{8} + \dots = 1$$



$$r + r(1-r) + r(1-r)^2 + \dots = 1$$


Geometric Series I

$$\sum_{n=0}^{\infty} ar^n = \frac{a}{1-r}$$



$$\frac{a + ar + ar^2 + ar^3 + \dots}{1/r} = \frac{ar}{1-r}$$

Geometric Series II




$$\Delta PQR \approx \Delta TSP$$


$$\therefore 1 + r + r^2 + \dots = \frac{1}{1-r}.$$

—Benjamin G. Klein and Irl C. Bivens

Geometric Series III



$$\frac{1}{4} + \frac{1}{16} + \frac{1}{64} + \frac{1}{256} + \dots = \frac{1}{3}$$




$$(1-r)^2 + r^2(1-r)^2 + r^4(1-r)^2 + \dots = \frac{(1-r)^2}{(1-r)^2 + 2r(1-r)} = \frac{1-r}{1+r}$$

$$1 + r^2 + r^4 + \dots = \frac{1}{1-r^2}$$

$$a + ar + ar^2 + \dots = \frac{a}{1-r}$$


Geometric Series IV



$$2 \left(\frac{1}{3} + 3 \cdot \frac{1}{27} + 9 \cdot \frac{1}{243} + \dots \right) = 1$$


$$2 \sum_{n=1}^{\infty} \frac{1}{3^n} = 1$$

$$\sum_{n=1}^{\infty} \frac{1}{3^n} = \frac{1}{2}$$



$$3 \sum_{n=1}^{\infty} \frac{1}{4^n} = 1$$

$$\sum_{n=1}^{\infty} \frac{1}{4^n} = \frac{1}{3}$$




$$4 \sum_{n=1}^{\infty} \frac{1}{5^n} = 1$$

$$\sum_{n=1}^{\infty} \frac{1}{5^n} = \frac{1}{4}$$

—Elizabeth M. Markham

Gabriel's Staircase


$$\sum_{k=1}^{\infty} kr^k = \frac{r}{(1-r)^2} \text{ for } 0 < r < 1$$




$$\sum_{k=1}^{\infty} kr^k = \sum_{k=1}^{\infty} \sum_{i=k}^{\infty} r^i = \frac{r}{(1-r)^2}$$

Differentiated Geometric Series


$$1 + 2\left(\frac{1}{2}\right) + 3\left(\frac{1}{4}\right) + 4\left(\frac{1}{8}\right) + \dots = 4$$



$$1 + 2r + 3r^2 + 4r^3 + \dots = \left(\frac{1}{1-r}\right)^2, \quad 0 \leq r < 1$$




$$\frac{1}{1 \cdot 2} + \frac{1}{2 \cdot 3} + \cdots + \frac{1}{n(n+1)} = \frac{n}{n+1}$$




—Roman W. Wong

The Series of Reciprocals of Triangular Numbers

$$\frac{1}{1} + \frac{1}{3} + \frac{1}{6} + \cdots + \frac{1}{\binom{n+1}{2}} + \cdots = 2$$



The Alternating Harmonic Series



$$\frac{1}{2} \left(\frac{2}{3} - \frac{2}{4} \right) = \frac{1}{3} - \frac{1}{4};$$


$$\frac{1}{4} \left(\frac{4}{5} - \frac{4}{6} \right) = \frac{1}{5} - \frac{1}{6}, \quad \frac{1}{4} \left(\frac{4}{7} - \frac{4}{8} \right) = \frac{1}{7} - \frac{1}{8};$$

$$\frac{1}{2^n} \left(\frac{2^n}{2^n + 2k-1} - \frac{2^n}{2^n + 2k} \right) = \frac{1}{2^n + 2k-1} - \frac{1}{2^n + 2k}, \quad k = 1, 2, \dots, 2^{n-1}; \\ n = 1, 2, \dots.$$


$$\ln 2 = \int_1^2 \frac{dx}{x} = \left(1 - \frac{1}{2} \right) + \left(\frac{1}{3} - \frac{1}{4} \right) + \dots = 1 - \frac{1}{2} + \frac{1}{3} - \frac{1}{4} + \dots$$

—Mark Finkelstein

$$\sin(2n+1)\theta = \sin\theta + 2\sin\theta \sum_{k=1}^n \cos 2k\theta$$



An Arctangent Identity and Series



$$\arctan n + \arctan \frac{1}{n^2+n+1} = \arctan(n+1)$$


$$\arctan \frac{1}{n^2+n+1} = \arctan(n+1) - \arctan n$$

$$\sum_{n=0}^{\infty} \arctan \frac{1}{n^2+n+1} = \lim_{N \rightarrow \infty} \arctan(N+1) = \frac{\pi}{2}$$

Miscellaneous

A 2×2 Determinant is the Area of a Parallelogram	133
Area of the Parallelogram Determined by Vectors (a,b) and (c,d)	134
The Characteristic Polynomials of AB and BA are Equal	135
The Gaussian Quadrature as the Area of Either Trapezoid	136
Inductive Construction of an Infinite Chessboard with Maximal Placement of Nonattacking Queens	137
Combinatorial Identities	138
$3 \sum_{j=0}^n \binom{3n}{3j} = 8^n + 2(-1)^n$, by Inclusion-Exclusion in Pascal's Triangle	139
The Existence of Infinitely Many Primitive Pythagorean Triples	140
Pythagorean Triples via Double Angle Formulas	141
The Problem of the Calissons	142
Recursion	143
$\prod_{k=1}^n k^k \cdot (k!) = (n!)^{n+1}$	144

A 2×2 Determinant is the Area of a Parallelogram







$$\begin{vmatrix} a & b \\ c & d \end{vmatrix} = ad - bc = \left\| \begin{array}{|c|c|} \hline \square & \square \\ \hline \end{array} \right\| - \left\| \begin{array}{|c|c|} \hline \square & \square \\ \hline \end{array} \right\| = \left\| \begin{array}{|c|c|} \hline \diagup & \diagdown \\ \hline \end{array} \right\|$$

—Solomon W. Golomb

Area of the Parallelogram Determined by

$$\text{Vectors } (a,b) \text{ and } (c,d) = \pm \begin{vmatrix} a & b \\ c & d \end{vmatrix} = \pm (ad - bc)$$




The Characteristic Polynomials of AB and BA are Equal

$$-\lambda^n |AB - \lambda I| = \left| \begin{pmatrix} A & AB - \lambda I \\ \lambda I & 0 \end{pmatrix} \right| = \left| \begin{pmatrix} A & I \\ \lambda I & B \end{pmatrix} \begin{pmatrix} I & B \\ 0 & -\lambda I \end{pmatrix} \right| = \left| \begin{pmatrix} A & I \\ \lambda I & B \end{pmatrix} \right| (-\lambda)^n$$

$$-\lambda^n |BA - \lambda I| = \left| \begin{pmatrix} 0 & \lambda I \\ BA - \lambda I & \lambda B \end{pmatrix} \right| = \left| \begin{pmatrix} A & I \\ \lambda I & B \end{pmatrix} \begin{pmatrix} -I & 0 \\ A & \lambda I \end{pmatrix} \right| = \left| \begin{pmatrix} A & I \\ \lambda I & B \end{pmatrix} \right| (-\lambda)^n$$


The Gaussian Quadrature as the Area of Either Trapezoid

$$\frac{1}{2}(b - a)(f(\bar{a}) + f(\bar{b})) = \frac{1}{2}(b - a)(h(a) + h(b))$$



—Mike Akerman

Inductive Construction of an Infinite Chessboard with Maximal Placement of Nonattacking Queens



REFERENCES


1. Dean S. Clark and Oved Shisha, Invulnerable Queens on an Infinite Chessboard, *Annals of the New York Academy of Sciences, The Third International Conference on Combinatorial Mathematics*, 1989, 133-139.
2. M. Kraitchik, *La Mathématique des Jeux ou Récréations Mathématiques*, Imprimerie Stevens Frères, Bruxelles, 1930, 349-353.

—Dean S. Clark and Oved Shisha

Combinatorial Identities


$$\binom{n}{2} = \frac{1}{2}(n^2 - n) = \sum_{i=1}^{n-1} i$$

$$\binom{n+1}{2} = \binom{n}{2} + n$$




—James O. Chilaka

$3 \sum_{j=0}^n \binom{3n}{3j} = 8^n + 2(-1)^n$, by Inclusion-Exclusion in Pascal's Triangle



$$\sum_{j=0}^n \binom{3n}{3j} = \sum_{j=1}^{3n-1} (-1)^{j-1} 2^{3n-j} = -2^{3n} \sum_{j=1}^{3n-1} \left(-\frac{1}{2}\right)^j = \frac{8^n + 2(-1)^n}{3}.$$


The Existence of Infinitely Many Primitive Pythagorean Triples



$$n^2 = 2k+1 \Rightarrow k^2 + n^2 = (k+1)^2 \quad \& \quad (k, k+1) = 1$$

—Charles Vanden Eynden

Pythagorean Triples via Double Angle Formulas




$$\begin{aligned}m &> n > 0 \\ m, n &\in \mathbb{I}\end{aligned}$$

$$\left\{ \begin{array}{l} \sin \theta = \frac{n}{\sqrt{m^2 + n^2}} \\ \cos \theta = \frac{m}{\sqrt{m^2 + n^2}} \end{array} \right.$$


$$\sin 2\theta = \frac{2mn}{m^2 + n^2}$$

$$\cos 2\theta = \frac{m^2 - n^2}{m^2 + n^2}$$



The Problem of the Calissons



A *calisson* is a French sweet that looks like two equilateral triangles meeting along an edge. Calissons could come in a box shaped like a regular hexagon, and their packing would suggest an interesting combinatorial problem. Suppose a box with side of length n is filled with sweets of sides of length 1. The short diagonal of each calisson in the box is parallel to a pair of sides of the box.



We refer to these three possibilities by saying that a calisson admits three distinct orientations.


THEOREM: *In any packing, the number of calissons with a given orientation is one-third of the total number of calissons in the box.*

PROOF:




—Guy David and Carlos Tomei


Recursion




$$A_2 = 3$$



$$A_3 = 2A_2 + 1$$



$$A_4 = 2A_3 + 1$$




$$A_5 = 2A_4 + 1$$

$$A_2 = 3 \text{ } \& \text{ } A_n = 2A_{n-1} + 1 \Leftrightarrow A_n = 2(2^{n-1}) - 1 = 2^n - 1$$

—Shirley Wakin

$$\prod_{k=1}^n k^k \cdot k! = (n!)^{n+1}$$



—Edward T. H. Wang

Sources

page	source
Geometry & Algebra	
3	Howard Eves, <i>Great Moments in Mathematics (Before 1650)</i> , The Mathematical Association of America, Washington, 1980, pp. 27-28.
4	Howard Eves, <i>Great Moments in Mathematics (Before 1650)</i> , The Mathematical Association of America, Washington, 1980, pp. 29-32.
5	Howard Eves, <i>Great Moments in Mathematics (Before 1650)</i> , The Mathematical Association of America, Washington, 1980, pp. 31, 33.
6	Howard Eves, <i>Great Moments in Mathematics (Before 1650)</i> , The Mathematical Association of America, Washington, 1980, pp. 29-30.
7	Howard Eves, <i>Great Moments in Mathematics (Before 1650)</i> , The Mathematical Association of America, Washington, 1980, pp. 34-36.
8	<i>College Mathematics Journal</i> , vol. 17, no. 5 (Nov. 1986), p. 422.
9	<i>College Mathematics Journal</i> , vol. 20, no. 1 (Jan. 1989), p. 58.
10	<i>Mathematics Magazine</i> , vol. 50, no. 3 (May 1977), p. 162.
11	<i>Mathematics Magazine</i> , vol. 48, no. 4 (Sept.-Oct. 1975), p. 198.
12	<i>College Mathematics Journal</i> , vol. 21, no. 5 (Nov. 1990), p. 393.
13	<i>Mathematics Magazine</i> , vol. 67, no. 5 (Dec. 1994), p. 354.
14	<i>College Mathematics Journal</i> , vol. 17, no. 4 (Sept. 1986), p. 338.
15	<i>Mathematics Magazine</i> , vol. 62, no. 3 (June 1989), p. 190.
16	<i>College Mathematics Journal</i> , vol. 22, no. 5 (Nov. 1991), p. 420.
17	<i>Mathematics Magazine</i> , vol. 66, no. 1 (Feb. 1993), p. 13.
18	<i>Mathematics Magazine</i> , vol. 65, no. 5 (Dec. 1992), p. 356.
19	<i>Mathematics Magazine</i> , vol. 56, no. 2 (March 1983), p. 110.
20	<i>Mathematics Magazine</i> , vol. 57, no. 4 (Sept. 1984), p. 231.
21	<i>Mathematics Magazine</i> , vol. 64, no. 2 (April 1991), p. 138.
22	<i>Mathematics Magazine</i> , vol. 66, no. 3 (June 1993), p. 180.
23	<i>Mathematics Magazine</i> , vol. 64, no. 2 (April 1991), p. 114.
24	<i>Mathematics Magazine</i> , vol. 68, no. 2 (April 1995), p. 109.
25	<i>Mathematics Magazine</i> , vol. 67, no. 4 (Oct. 1994), p. 302.

pagesource**Trigonometry, Calculus & Analytic Geometry**

- 29 *Mathematics Magazine*, vol. 64, no. 2 (April 1991), p. 97.
 30 *Mathematics Magazine*, vol. 62, no. 5 (Dec. 1989), p. 317.
 31 *Mathematics Magazine*, vol. 61, no. 4 (Oct. 1988), p. 259.
 32 *Mathematics Magazine*, vol. 63, no. 5 (Dec. 1990), p. 342.
 33 *Mathematics Magazine*, vol. 64, no. 2 (April 1992), p. 103.
 34 *College Mathematics Journal*, vol. 20, no. 1 (Jan. 1989), p. 51.
 35 *American Mathematical Monthly*, vol. 49, no. 5 (May 1942), p. 325.
 36 *Mathematics Magazine*, vol. 61, no. 5 (Dec. 1988), p. 281.
 37 *Mathematics Magazine*, vol. 61, no. 2 (April 1988), p. 113.
 38 *Mathematics Magazine*, vol. 62, no. 4 (Oct. 1989), p. 267.
 39 *College Mathematics Journal*, vol. 18, no. 2 (March 1987), p. 141.
 40 *Mathematics Magazine*, vol. 42, no. 1 (Jan.-Feb. 1969), pp. 40-41.
 41 *College Mathematics Journal*, vol. 16, no. 1 (Jan. 1985), p. 56.
 42 Richard Courant, *Differential and Integral Calculus*, p. 219,
copyright © 1937, Interscience. Reprinted by permission of John
Wiley & Sons, Inc.
 43 *College Mathematics Journal*, vol. 18, no. 1 (Jan. 1987), p. 52.
 44 *Mathematics Magazine*, vol. 64, no. 3 (June 1991), p. 175.
 45 *Mathematics Magazine*, vol. 66, no. 1 (Feb. 1993), p. 39.

Inequalities

- 49 *Mathematics Magazine*, vol. 50, no. 2 (March 1977), p. 98.
 50 *Mathematics Magazine*, vol. 59, no. 1 (Feb. 1986), p. 11.
 51 *College Mathematics Journal*, vol. 16, no. 3 (June 1985), p. 208.
 52 *College Mathematics Journal*, vol. 19, no. 4 (Sept. 1988), p. 347.
 53 *Mathematics Magazine*, vol. 60, no. 3 (June 1987), p. 158.
 54 *College Mathematics Journal*, vol. 21, no. 3 (May 1990), p. 227.
 55 *College Mathematics Journal*, vol. 20, no. 3 (May 1989), p. 231.
 56 *Mathematics Magazine*, vol. 69, no. 1 (Feb. 1996), p. 64.
 58 *Mathematics Magazine*, vol. 60, no. 3 (June 1987), p. 165.
 59 *Mathematics Magazine*, vol. 64, no. 1 (Feb. 1991), p. 31.
 60 *Mathematics Magazine*, vol. 63, no. 3 (June 1990), p. 172.
 61 I. Reprinted by permission from *Mathematics Teacher*, vol. 81,
no. 1 (Jan. 1988), p. 63, author Li Changming, copyright © 1988 by
the National Council of Teachers of Mathematics, Inc.

page	source
Inequalities (continued)	
61	<i>II. Mathematics Magazine</i> , vol. 67, no. 1 (Feb. 1994), p. 34.
62	<i>Mathematics Magazine</i> , vol. 67, no. 5 (Dec. 1994), p. 374.
63	<i>Mathematics Magazine</i> , vol. 66, no. 1 (Feb. 1993), p. 65.
64	<i>Mathematics Magazine</i> , vol. 67, no. 1 (Feb. 1994), p. 20.
65	<i>Mathematics Magazine</i> , in press.
66	<i>College Mathematics Journal</i> , vol. 24, no. 2 (March 1993). p. 165.

Integer Sums

- 69 *Scientific American*, vol. 229, no. 4 (Oct. 1973), p. 114.
- 70 *Mathematics Magazine*, vol. 57, no. 2 (March 1984), p. 104.
- 71 Reprinted by permission from *Historical Topics for the Mathematics Classroom*, p. 54, author Bernard H. Gundlach, copyright © 1969 by the National Council of Teachers of Mathematics, Inc.
- 73 *Mathematics Magazine*, vol. 64, no. 2 (April 1991), p. 103.
- 74 *Scientific American*, vol. 229, no. 4 (Oct. 1973), p. 115.
- 75 *Mathematics Magazine*, vol. 66, no. 3 (June 1993), p. 166.
- 76 *Mathematics Magazine*, vol. 59, no. 2 (April 1986), p. 92.
- 77 *Mathematics Magazine*, vol. 57, no. 2 (March 1984), p. 92.
- 78 *Scientific American*, vol. 229, no. 4 (Oct. 1973), p. 115.
College Mathematics Journal, vol. 22, no. 2 (March 1991), p. 124.
- 79 *College Mathematics Journal*, vol. 20, no. 3 (May 1989), p. 205.
- 80 *Mathematics Magazine*, vol. 56, no. 2 (March 1983), p. 90.
- 81 *College Mathematics Journal*, vol. 20, no. 2 (March 1989), p. 123.
- 82 I. *Mathematics Magazine*, vol. 60, no. 5 (Dec. 1987), p. 291.
 II. *Mathematics Magazine*, vol. 65, no. 2 (April 1992), p. 90.
- 83 M. Bicknell & V. E. Hoggatt, Jr. (eds.), *A Primer for the Fibonacci Numbers*, The Fibonacci Association, San Jose, 1972, p. 147.
- 84 *Mathematical Gazette*, vol. 49, no. 368 (May 1965), p. 199.
- 85 *Mathematics Magazine*, vol. 50, no. 2 (March 1977), p. 74.
- 86 *Mathematics Magazine*, vol. 58, no. 1 (Jan. 1985), p. 11.
- 87 *Mathematics Magazine*, vol. 62, no. 4 (Oct. 1989), p. 259.
Mathematical Gazette, vol. 49, no. 368 (May 1965), p. 200.
- 88 *Mathematics Magazine*, vol. 63, no. 3 (June 1990), p. 178.
- 89 *Mathematics Magazine*, vol. 62, no. 5 (Dec. 1989), p. 323.
- 90 *Mathematics Magazine*, vol. 65, no. 3 (June 1992), p. 185.

pagesource**Integer Sums (continued)**

- 91 *Mathematics Magazine*, vol. 68, no. 5 (Dec. 1995), p. 371.
 92 *Mathematics Magazine*, vol. 65, no. 1 (Feb. 1992), p. 55.
 93 *Mathematics Magazine*, vol. 66, no. 5 (Dec. 1993), p. 329.
 94 *Mathematics Magazine*, vol. 63, no. 5 (Dec. 1990), p. 314.
 95 *College Mathematics Journal*, vol. 23, no. 5 (Nov. 1992), p. 417.
 96 Richard K. Guy, written communication.
 97 *Mathematics Magazine*, vol. 62, no. 1 (Feb. 1989), p. 27.
 98 *Mathematics Magazine*, vol. 62, no. 2 (April 1989), p. 96.
 99 *College Mathematics Journal*, vol. 18, no. 4 (Sept. 1987), p. 318.
 100 *Mathematics Magazine*, vol. 66, no. 5 (Dec. 1993), p. 325.
 101 *Mathematics Magazine*, vol. 58, no. 4 (Sept. 1985), pp. 203, 236.
 102 *Mathematics Magazine*, vol. 63, no. 1 (Feb. 1990), p. 25.
 103 *Mathematics Magazine*, vol. 55, no. 2 (March 1982), p. 97.
 104 Richard K. Guy, written communication.
 105 *Mathematics Magazine*, vol. 67, no. 4 (Oct. 1994), p. 293.
 106 Richard K. Guy, written communication.
 107 *Mathematics Magazine*, vol. 60, no. 5 (Dec. 1987), p. 327.
 108 *Mathematics Magazine*, vol. 63, no. 4 (Oct. 1990), p. 225.
 109 *American Mathematical Monthly*, vol. 95, no. 8 (Oct. 1988), pp. 701, 709.
 110 *Mathematics Magazine*, vol. 66, no. 5 (Dec. 1993), p. 316.
 111 *Mathematics Magazine*, vol. 63, no. 5 (Dec. 1990), p. 349.

Sequences & Series

- 115 *Mathematics Magazine*, vol. 68, no. 1 (Feb. 1995), p. 41.
 116 *Mathematics Magazine*, vol. 67, no. 5 (Dec. 1994), p. 379.
 117 *Mathematics Magazine*, vol. 66, no. 3 (June 1993), p. 179.
 118 *Mathematics Magazine*, vol. 54, no. 4 (Sept. 1981), p. 201.
 119 *Mathematics Magazine*, vol. 60, no. 3 (June 1987), p. 177.
 120 *Mathematics Magazine*, vol. 61, no. 4 (Oct. 1988), p. 219.
 121 *Mathematics Magazine*, vol. 67, no. 3 (June 1994), p. 230.
 122 *Mathematics Magazine*, vol. 66, no. 4 (Oct. 1993), p. 242.
 123 *Mathematics Magazine*, vol. 67, no. 3 (June 1994), p. 209.
 124 *Mathematics Magazine*, vol. 62, no. 5 (Dec. 1989), pp. 332-333.
 126 *Mathematics Magazine*, vol. 65, no. 5 (Dec. 1992), p. 338.
 127 *Mathematics Magazine*, vol. 64, no. 3 (June 1991), p. 167.

page	source
------	--------

Sequences & Series (continued)

- 128 *American Mathematical Monthly*, vol. 94, no. 6 (June-July 1988), pp. 541-42.
- 129 *Mathematics Magazine*, vol. 65, no. 2 (April 1992), p. 136.
- 130 *Mathematics Magazine*, vol. 64, no. 4 (Oct. 1991), p. 241.

Miscellaneous

- 133 *Mathematics Magazine*, vol. 58, no. 2 (March 1985), p. 107.
- 134 *Mathematics Magazine*, vol. 64, no. 5 (Dec. 1991), p. 339.
- 135 *Mathematics Magazine*, vol. 61, no. 5 (Dec. 1988), p. 294.
- 136 *Mathematics Magazine*, vol. 60, no. 2 (April 1987), p. 89.
- 137 *Mathematics Magazine*, vol. 61, no. 2 (April 1988), p. 98.
- 138 *Mathematics Magazine*, vol. 52, no. 4 (Sept. 1979), p. 206.
- 139 *Mathematics Magazine*, vol. 63, no. 1 (Feb. 1990), p. 29.
- 140 *Elementary Number Theory*, Random House, New York, 1987, p. 227.
- 141 *Mathematics Magazine*, vol. 67, no. 3 (June 1994), p. 187.
- 142 *American Mathematical Monthly*, vol. 96, no. 5 (May 1989), pp. 429-30.
- 143 *Mathematics Magazine*, vol. 62, no. 3 (June 1989), p. 172.
- 144 *College Mathematics Journal*, vol. 20, no. 2 (March 1989), p. 152.

Index of Names

- Ajose, Sunday A. 121
Akerman, Mike 136
Amir-Mo  z, Ali R. 99
Aristarchus of Samos 63
Ayoub, Ayoub B. 43, 44
- Beekman, Richard M. 45
Bernoulli, Jacques 65
Bh  skara .4
Bicknell, M. 147
Bivens, Irl C. 120
Bronson, Robert 111
Brousseau, Alfred 83
Brueningsen, Christopher 111
Burk, Frank 41
- Cauchy, Augustin-Louis 64
Cavalieri, Bonaventura 25
Chilaka, James O. 76, 80, 138
Chuang, Pi-Chun 81
Chuquet, Nicolas 61
Clark, Dean S. 137, 139
Coble, Scott 13
Courant, Richard 42
Cupillari, Antonella 87
- David, Guy 142
DeKleine, H. Arthur 36
Dence, Thomas P. 117
Diophantus of Alexandria 22
Drude, K. Ann 21
Dudeney, H. E. 6
- Eddy, Roland H. 16, 18, 51
Eisenman, R. L. 40
Elsner, Thomas 10
Euclid 5
Eves, Howard 145
- Fibonacci (Leonardo of Pisa) 83
Finkelstein, Mark 128
Fisher, J. Chris 129
Fry, Alan L. 86
- Galileo Galilei 115
Gallant, Charles D. 19, 49, 59
Gardner, Martin 69, 74, 78
Garfield, James A. 7
Gau, Yihnan David 134
Gentile, Enzo R. 9
Gibbs, Richard A. 60
Golomb, Solomon W. 84, 133
Gundlach, Bernard H. 147
Guy, Richard 148
- Hardy, Michael 8
Harris, Edward M. 39
Hoggatt, V. E. 147
Houston, David 141
- Isaacs, Rufus 11
- Kalman, Dan 78
Kim, Hee Sik 75
Kin  canon, Eric 12
Klein, Benjamin G. 120
Koh, E. L. 129
Kung, Sidney H. 17, 25, 29, 30,
32, 33, 54, 79, 98, 135
- Landauer, Edwin G. 101
Lehel, Jen   73
Li Changming 61
Logothetti, Dave 23, 82
Love, J. Barry 85
Lushbaugh, Warren 87


Markham, Elizabeth M.	92, 122	Schwarz, Hermann A.	64
Miller, William A.	100	Shisha, Oved	137
Mollweide, Karl	36	Sipka, Timothy A.	31
Montucci, Paolo	52	Siu, Man-Keung	77
Nair, N. Gopalakrishnan	93	Snover, Steven L.	82
Nakhli, Fouad	14, 58	Swain, Stuart G.	123
Napier, John	66		
Nicomachus of Gerasa	71	Tomei, Carlos	142
Page, Warren	52, 103, 118	Vanden Eynden, Charles	140
Pascal, Blaise	139	Viviani, Vincenzo	15
Pooley, Sam	21		
Pouryoussefi, Farhood	89	Wakin, Shirley	20, 107, 143
Ptolemy, Claudius	33	Wang, Edward T. H.	144
Pythagoras	3-9, 140, 141	Walker, R. J.	35
Richards, Ian	70	Webb, J. H.	119
Romaine, William	37	Wolf, Samuel	15
Schattschneider, Doris	50	Wong, Roman W.	126
Schrage, Georg	90	Wu, T. C.	97
		Zerger, Monte J.	91, 94

Technical Note

The manuscript for this book was edited and printed using Microsoft® Word 5.1 on an Apple® Macintosh™ IIIfx computer. The graphics were produced in Claris™ MacDraw® Pro 1.5v1. The text is set in the Palatino font, with special characters in the Symbol font. Many of the displayed equations were produced with MacΣqn™ 3.0 from Software for Recognition Technologies. The manuscript was printed on an Apple® Laserwriter® II NTX.

PROOFS WITHOUT WORDS

EXERCISES IN VISUAL THINKING



Roger Nelsen received his Ph.D. in Mathematics from Duke University. Since 1969 he has taught at Lewis and Clark College, in Portland Oregon, where he is a professor of mathematics. He is currently an Associate Editor of the "Problems and Solutions" section of the *College Mathematics Journal*.

Journal. His expository and research papers in mathematics have appeared in the *American Mathematical Monthly*, *Mathematics Magazine*, *The College Mathematics Journal*, *Nature*, *Journal of Applied Probability*, *Communications in Statistics*, *Probability Theory and Related Fields*, *Statistics and Probability Letters*, *Sankhyā*, and the *Journal of Nonparametric Statistics*. His main research interests are in probability and mathematical statistics.

Just what are "proofs without words?" First of all, most mathematicians would agree that they certainly are not "proofs" in the formal sense. Indeed, the question does not have a simple answer. But, as you will see in this book, proofs without words are generally pictures or diagrams that help the reader see why a particular mathematical statement is true, and also to see how one could begin to go about proving it true. While in some proofs without words an equation or two may appear to help guide that process, the emphasis is clearly on providing visual clues to stimulate mathematical thought. Proofs without words bear witness to the observation that often in the English language to see means to understand, as in "to see the point of an argument."

Proofs without words have a long history. In this collection you will find modern renditions of proofs without words from ancient China, classical Greece, twelfth-century India—even one based on a published proof by a former President of the United States! However, most of the proofs are relatively more recent creations, and many are taken from the pages of MAA journals.

The proofs in this collection are arranged by topic into six chapters: Geometry and Algebra; Trigonometry, Calculus and Analytic Geometry; Inequalities; Integer Sums; Sequences and Series, and Miscellaneous. Teachers will find that many of the proofs without words in this collection are well suited for classroom discussion and for helping students to think visually in mathematics.

